

This photograph of the moon's southeast quadrant, showing the craters Tycho and Clavius, was made by the 3.5-inch Questar telescope, which weighs 7 pounds. It was made on 35-mm. film at sea level. There is remarkable detail, as much as found in pictures taken with much larger instruments. Compare it, for example, with Anton Kutter's fine 12-inch Schiefspiegler photograph on page 67 of the December, 1958, issue of *Sky and Telescope*.

Of course, large telescopes suffer much more from inferior seeing conditions than do small ones, and Questar sometimes outperforms instruments several times its size. Questar's highly corrected optical system can always be used at full aperture, while big telescopes of other types may have to be stopped down. The old statement that resolving power is proportional to aperture is not always true during actual observing at night.

Dr. G. P. Kuiper in his magnificent Photographic Lunar Atlas, which contains some of the finest moon plates ever taken, shows the cleft system west of Triesnecker on five pictures. The best resolution obtained is about 0".4, he says, matching the visual resolving power of an 11-inch telescope. Plate C4-e, by the 100-inch reflector at Mount Wilson, is an unparalleled picture of this intricate cleft system. Horace Dall's 15.5-inch caught much of it in a picture on page 62 of Henry Paul's "Outer Space Photography." The miniscule Questar has also succeeded in photographing the chief clefts, as shown on page 14 of our latest booklet.

May we send you one?

**QUESTAR**  
NEW HOPE, PENNSYLVANIA

STAR-TESTED QUESTARS ARE USUALLY IN STOCK READY FOR IMMEDIATE DELIVERY, FROM \$995. TERMS ARE AVAILABLE. LET US SEND YOU THE QUESTAR BOOKLET.

## CONTENTS

**COVER:** Observing at the Cassegrainian focus of Haute Provence Observatory's main telescope. A rising section of floor makes access to the eyepiece easy; a heat shield and fans prevent distortions of the mirror because of the observer's proximity. When the Newtonian focus is used, a counterweight replaces the mirror backing. (See page 4.)

SOLAR MAGNETISM AND SUNSPOT FORMATION .....	3
FRANCE'S HAUTE PROVENCE OBSERVATORY	
— Jean Dufay and Charles Fehrenbach .....	4
RADIO URSIGRAMS	
— Martin H. Potter .....	12
E. E. BARNARD AND MILKY WAY PHOTOGRAPHY	
— Otto Struve .....	14
JOHANN KEPLER AND THE LAWS OF PLANETARY MOTION	
— Rufus Suter .....	22
ASTRONOMERS DISCUSS PROJECT WEST FORD .....	25
TEN YEARS OF SOLAR ECLIPSES .....	29
AMATEUR ASTRONOMERS .....	26
An Active Amateur in St. Paul	
BOOKS AND THE SKY .....	39
Planets, Stars, and Galaxies	
Captured Stars	
Radio Astronomy	
Rival Theories of Cosmology	
Handbuch fur Sternfreunde	
CELESTIAL CALENDAR .....	54
An Early-Morning Occultation of Aldebaran	
GETTING ACQUAINTED WITH ASTRONOMY .....	19
Jupiter's Satellites	
GLEANINGS FOR ATM's .....	47
An 8-Inch "Beavertail" Dall-Kirkham Reflector	
Checking Concentricity of Maksutov Correcting Lenses	
LETTERS .....	21
NEWS NOTES .....	10
OBSERVER'S PAGE .....	32
Mutual Phenomena of Jupiter's Satellites	
Observing the Moon — Lassell	
Deep-Sky Wonders	
OBSERVING THE SATELLITES .....	23
QUESTIONS .....	13
STARS FOR JULY .....	57

SKY AND TELESCOPE is published monthly by Sky Publishing Corporation, Harvard College Observatory, Cambridge 38, Mass. Second-class postage paid at Boston, Mass.

Subscription: \$5.00 per year in the United States and possessions; \$9.00 for two years; \$13.00 for three years. To Canada, Mexico, and all countries of the Pan American Postal Union, \$6.00 for one year; \$11.00 for two years; \$16.00 for three years. To all other foreign countries, \$7.00 for one year; \$13.00 for two years; \$19.00 for three years. Canadian and other foreign remittances should be made in United States currency. Single copy 50 cents. Back issues: November, 1958, to present, 50 cents per copy. Vols. I through XVI (November, 1941, to October, 1957), order from Walter J. Johnson, Inc., 111 Fifth Ave., New York 3, N. Y.

All notices of change of address must be sent one month in advance and accompanied by old and new addresses. When sending your renewal order, or writing in regard to your subscription, your current mailing address must be given. Return our bill form with renewal payment. Circulation manager, Nancy R. Bolton. Office staff: Sarah E. Bulger, Virginia K. Cox, J. Caroline Nason, Mary Silva, John Simmons, Helen B. Sniegiecki, Stacia Strong, Aubrey S. Wheaton, Jr.

Editorial and advertising offices: 49-50-51 Bay State Rd., Cambridge 38, Mass. Phone UN 4-7360. Assistant editors: Gail B. Carney, advertising; William E. Shawcross, departments; Raymond N. Watts, Jr., features. Assistants: George A. Blair, Willen P. Schrauwen. Advertising rates are listed in STANDARD RATE AND DATA SERVICE, or sent upon request; closing date (except covers) is 20th of second month preceding date of issue. Unsolicited articles and pictures are welcome, but prompt attention and return are not guaranteed.

The principal articles are indexed in THE READERS' GUIDE TO PERIODICAL LITERATURE. Although the contents of this magazine, its text, pictures, diagrams, and charts, are copyrighted with all rights reserved, we are pleased to give permission, upon request, for the use of SKY AND TELESCOPE material for educational and scientific purposes, with the exception of the monthly star and planet charts, which may not be copied or reproduced in any manner for any purpose. Permission to use matter from signed articles, and photographs and diagrams for which credit is given, must also be obtained from the person, observatory, institution, or agency concerned.

## Solar Magnetism and Sunspot Formation

**M**OUNT WILSON Observatory was founded more than half a century ago, especially for studies of the sun and sunspots. Now its assistant director, Horace W. Babcock, offers a far-reaching synthesis of current ideas and theories concerning the sun's magnetic field and the nature of sunspots. Although the latter have been among the most thoroughly observed astronomical phenomena, the mechanism of their formation has remained a mystery.


Combining his own studies of solar magnetism with those of his father, Harold D. Babcock, and the theoretical proposals of many other astronomers, Dr. Babcock offers the first comprehensive explanation of why sunspots vary in frequency, size, and latitude, why the magnetic fields of sunspot groups and of the entire sun reverse their polarities every 11 years, and why there are variations in the numbers and sizes of clouds of hot gas that are ejected from the sun and float through the solar system. The Babcocks have pioneered in developing and using special magnetographs for recording daily the details of magnetic flux in the photosphere (SKY AND TELESCOPE, October, 1954, page 423; September, 1958, page 555).

The new theory is based on the fact that all parts of the sun — a sphere of gas 865,000 miles in diameter — do not rotate together. Its equatorial belt rotates once every 25 days, while zones near the pole require 34 days to complete one turn. Thus, every three years the equatorial regions of the sun make five more rotations than sections in latitudes 50° north or south. Dr. Babcock believes that this wraps the lines of force of the main solar magnetic field, which are embedded in the electrically conducting gases of the sun's interior, in opposite directions in the northern and southern hemispheres.

Three years is about the time required for the magnetic wrappings to build up prior to the onset of a new sunspot cycle. As they are wound around the sun in a layer roughly 40,000 miles beneath the surface, the lines of force are stretched and tightened, and the sub-surface field becomes as much as 45 times stronger, growing most intense in the intermediate latitudes (about 30° north and south). New-cycle sunspots generally appear in these regions.

The comparatively fast-moving layers near the equator induce twisting of the lines of force to form what Dr. Babcock calls "magnetic ropes." In such a rope there may be local constrictions due to excess twisting, increasing its energy and making it unstable. This may manifest it-

(Continued on page 11)



A general view of the Haute Provence Observatory, showing from left to right an antenna of the radio interferometer, the dome of the 76-inch telescope, workshop, laboratories, the dome of the 47-inch, and living quarters.

## France's Haute Provence Observatory

JEAN DUFAY and CHARLES FEHRENBACH, *Haute Provence Observatory*

IN 1936, responding to a wish of French astronomers, Mme. Irène Joliot-Curie, first under-secretary of state for scientific research, decided to establish a well-equipped astrophysical observatory in a favorable climate. The Haute Provence


Observatory has since become one of the principal institutions operating directly under the National Center of Scientific Research, which is part of the French ministry of national education.

The site was quickly chosen, thanks to long reconnoitering previously organized by André Danjon, with the collaboration of André Couder and Jean Dufay. Five years of testing with a temporarily mounted 32-inch telescope at Forcalquier verified that the Basses-Alpes region of southeastern France was perfectly suited for large instruments.

After further detailed site testing, the observatory was constructed at St. Michel, on a rocky plateau 2,000 feet high, less than six miles from Forcalquier and about 60 miles north of Marseilles. This region of Haute Provence, on the last foothills of the Alps, possesses a typically Mediterranean climate, with very few rainy days. An average year has 250 nights when observing is possible, over 100 of which are clear throughout. The number of nights offering almost perfect seeing is also very satisfying.

Undertaken in the beginning of 1938, construction work proceeded very rapidly until the war, under the inspiration of

The dome of the 76-inch reflector, Europe's third largest telescope. The shutter moves up over the dome, and the leaves shown folded at the slit's bottom can be raised as a windscreen.







Inside the dome of the 76-inch reflector. An observer stands at the Newtonian focus near the tube's upper end. His observing platform rides up and down the heavy column (far right), whose base may be moved completely around the edge of the dome. The Newtonian secondary can be turned to reflect light to any one of four positions spaced around the tube, so the plateholder can be conveniently placed. In the lower right an assistant sits at the control console. The telescope may also be used as a Cassegrainian (front cover) or as a coude, whose spectrograph chamber is at the south end of the polar axis, beneath the console. All illustrations with this article are courtesy Haute Provence Observatory.



The main parts of the system for air-conditioning the interior of the 76-inch telescope are visible in this view. Protruding at the top is the porous octagonal inner casing. Fans at the tube's base maintain smooth air flow down the tube, and also help bring the mirror quickly into thermal equilibrium with the outside air. Two fans (see front cover) are placed back of the mirror when the Cassegrainian focus is used. One of the dome's fans appears behind the finder telescope at lower right in the picture.

the great physicist Jean Perrin, successor to Mme. Joliot-Curie. Little was done, of course, from 1940 to 1945; yet the first telescope, a 47-inch, began operation in 1943.

The observatory at present possesses four standard reflectors, with apertures of 24 inches (Cassegrainian), 32 inches (Newtonian and Cassegrainian), 47 inches (Newtonian), and 76 inches (Newtonian, Cassegrainian, and coude). It also has an  $f/2$  Schmidt camera with a correcting plate 12 inches in diameter.

For special work, there are two objective-prism spectrographs of the Fehrenbach type, with apertures of six and 16 inches; a Lyot coronagraph; and spectrographs and photometers for observing the


airglow of the night sky. The optical parts of the principal instruments, including the three largest reflectors, were figured by Dr. Couder at the Paris Observatory's optical laboratory. We have a large interferometer for the study of galactic and extragalactic radio sources. Especially important are our laboratory facilities and electrical and machine shops.

The Haute Provence Observatory is at the disposal of all French astrophysicists, who may reside on the grounds while using the facilities for specific projects. Accommodations include a dining room and a 30-room guest house, the Maison Jean Perrin. Observing is the only activity at St. Michel; guest scientists plan their programs and analyze data at their home observatories.


There are, therefore, no astronomers now in permanent residence at the observatory, although coauthor Fehrenbach has been going there for eight consecutive years, and a number of other researchers have worked there several years. On the other hand, an administrator and numerous technical personnel, including engineers, mechanics, electricians, and observing assistants, live permanently at St. Michel, in addition to the usual domestic staff. Administration of the observatory was originally undertaken by coauthor Dufay, director of Lyons Observatory. In 1943 he was joined by Fehrenbach, who is also director of Marseilles Observatory.

A directing committee, presided over by Dr. Danjon, meets regularly to plan the observing schedule and assign telescope time. Foreign astronomers are authorized to work at the observatory under the same conditions as our countrymen.

The largest telescope of Haute Provence Observatory was put in operation in July, 1958. Its primary mirror, used with an aperture of 75 inches, was cast in



At the coude focus is a grating spectrograph. As shown in the diagram above, the field of view is first oriented by a field rotator. The light then passes through a rocking plate that moves the image back and forth to widen the spectrum. The collimating mirror reflects a parallel beam back to the grating, which then throws the spectrum into one of the five cameras. For guiding, the observer views the star's image which is reflected from the polished jaws of the slit. Camera II, not shown, is behind I and III, and all three rotate into position on a single axis; tilting the grating brings the others into play. Below is a side view of the spectrograph, showing the light entrance, guiding eyepiece, and the housing for camera IV. The whole assembly moves on rails.





This vast bubble of glowing gas is the planetary nebula NGC 7635 in Cassiopeia, recorded in a one-hour exposure with the 76-inch reflector. With a maximum diameter of about 200 seconds of arc, the nebula appears as an 8th-magnitude object at right ascension  $23^h 18^m.5$ , declination  $+60^\circ 54'$  (1950 co-ordinates). South is at the top.

France in 1939, from low-expansion glass; but it could not be ground until after the war. It is relatively thin, weighing only 3,000 pounds; its cell has three fixed supports on the back and two on the side, aligning the optical axis. Thirty balanced supports compensate for flexure. The figure of this mirror is excellent, no aberration exceeding  $1/14$  wave. Quartz was used for the various secondary mirrors.

The telescope is pointed, with a precision of one minute of arc, by remote control from a console where the right ascension and declination of the observed object can be read directly. The Newtonian focus (focal length 31.5 feet,  $f/5$ ) is easily accessible by means of a movable observing platform, no matter where the


telescope is pointing in the sky. This platform rides up and down a tall column, which in turn stands on a movable ring-shaped portion of the floor.

Observing at the Cassegrainian focus (equivalent focal length 93.5 feet,  $f/15$ ) is facilitated by a rising floor 31.2 feet in diameter and having 11.8 feet of vertical

using the instrument's prime focus has been left open.


The dome, 66 feet in outer diameter, is situated above the third floor of a building of reinforced concrete, whose foundation is on bedrock. The north and south piers of the telescope have their own foundation, which anchors them solidly

**The sodium D lines of Alpha Cygni are widely spaced in this portion of a Haute Provence coude spectrogram.**





travel. At the coude focus (187 feet,  $f/30$ ), the light is conducted into a large laboratory where a constant temperature is maintained. Finally, the possibility of

together independently of the rest of the building. The first floor provides a spacious room where the primary mirror can be lowered in its cell by means of a



The region from 5150 to 5230 angstroms in the spectrum of Beta Pegasi, an M-type giant star, taken with Camera V of the coude spectrograph. Dark flutings are a titanium oxide band, converging leftward to a head at 5165 angstroms. This is enlarged about 10 times from a plate whose original scale was 4.1 angstroms per millimeter. The comparison spectra above and below are of an iron arc. A speck of dust on the slit caused the dark horizontal line spanning the star spectrum.



Two pictures of the same star field near Delta Orionis: left, a direct exposure, and right, an objective-prism photograph taken with the 6-inch camera shown at bottom right on this page. Stars are labeled by their Henry Draper Catalogue numbers and by their spectral types. The A- and B-type spectra mainly show dark lines of hydrogen; the G5 and K5 stars have many metallic lines. The spectrum plate is an 80-minute exposure; north is to the right, east above.

traveling crane, to be aluminized in a 7½-foot stainless-steel tank.

Particular precautions are taken to reduce temperature fluctuations in the dome's interior and to avoid air turbulence. The dome covering consists of two sheets of aluminum mounted on steel ribs and separated from each other by an insulating air space filled with crumpled aluminum foil.

It is desirable that the rays of light entering the dome and telescope travel within a laminar flow of air, so that no turbulence will exist along the light path. The observing slit can be partially closed with movable shades, leaving an opening only eight feet square above the upper



end of the telescope. The tube is double-covered. Exterior air that enters it is drawn through the inner casing of porous nylon and blown out into the dome by six fans placed on the metallic outer casing of the tube not far from the primary mirror. Eight larger fans are mounted in the dome opposite the slit. The slight vacuum they create in the building causes a smooth inward flow in front of the telescope.

This system, due to Dr. Couder, has been found in careful tests to be very effective. Almost perfect images can be obtained, if external atmospheric turbulence is also rather small. Currently, observations at the coude focus are being

made of stellar images whose diameters are between 0.3 and 0.8 second of arc.

A large spectrograph of Fehrenbach design, constructed by the Society for Research and Studies in Optics and Related Sciences, is installed at the coude focus, on a pier rigidly connected to the telescope supports. The spectrograph can be moved out of the way very rapidly, though it weighs over two tons, to permit use of other observing equipment, for example the Lallemand electronic image converter.

The 8-inch collimating mirror is 14.8 feet in focal length. Two Bausch and Lomb gratings, six by eight inches, are instantly interchangeable from a control



Above: Mounted on a large pegboard are (top) the 6-inch objective-prism camera with its guiding telescope and, below them, a 12-inch Schmidt camera.

Left: The observatory's 24-inch reflector equipped with a Lallemand image converter, which greatly shortens exposure times.




panel at the top of the spectrograph, and send the light into any one of five cameras. The largest of these has a 31-inch mirror of 7.38-foot focus, and a photographic plate 14 inches long. With the first grating, having 1,200 lines per millimeter, it covers the spectral region from 3600 to 7200 angstroms in three sections. The dispersion is 4.1 angstroms per millimeter in the violet and visible regions, and 3.6 in the extreme red. This camera can record spectra of 8th-magnitude stars in a six-hour exposure. With the second grating (771 lines per millimeter), the same camera produces dispersions of 3.2 angstroms per millimeter in the violet and 6.4 in the near infrared.

The other cameras furnish dispersions in the neighborhood of 10, 20, 40, and 50 angstroms per millimeter. The two fastest cameras are  $f/1.1$ , and are particularly suited to the observation of nebulae.

More than a thousand plates have been made since the spectrograph was installed in July, 1959, and as the reproduced spectra show, definition is excellent.

Several other spectrographs have been planned for the Cassegrainian focus. A recently completed one is specially intended to measure radial velocities. Another will soon be installed for studying faint stars and nebulae in the visible and near-infrared spectral regions. Finally, an extremely fast spectrograph ( $f/0.47$ ) is planned to be used at the Newtonian focus.

Two novel Haute Provence telescopes are 6- and 16-inch refractors with objective prisms for the determination of radial velocities of stars. Their design was conceived by Fehrenbach and developed by Dr. Danjon. An objective-prism camera has the advantage of being able to photograph the spectra of many stars at once, but it is not possible to interpose a laboratory spectrum to furnish wave length



This instrument, designed by the French astronomer J. E. Blamont, is used for the study of free sodium in the upper atmosphere. The telescope receives radiation from atmospheric sodium, which is measured by the fluorescence it causes in a tube of heated sodium vapor within a magnetic field.

calibration. However, if two exposures are made, with the prism rotated 180 degrees between them, each star produces two spectra, one with blue to the left, the other with blue to the right.

The prisms used are compound ones that do not deviate light for a chosen wave length, this property holding over the entire field of the telescope. Hence the two spectra of a star will be symmetrical about this wave length, but displaced by an amount depending on the Doppler shift of the star, and therefore on its radial velocity. This method has proved very successful for the wholesale measure-

ment of the line-of-sight velocities of stars.


Other instruments at the observatory are of the conventional type. We hope to keep pace with new developments in instrumentation, so that French astrophysicists will have at St. Michel every facility to meet their observing requirements.

#### GEORGE A. DAVIS, JR., DIES

A widely known Buffalo, New York, astronomer died on June 4th at the age of 71. George A. Davis, Jr., was an authority on astronomical history and especially on star names, concerning which he contributed several articles to *SKY AND TELESCOPE*. He founded the Buffalo Astronomical Association in 1930, and was director of the Buffalo Museum of Science's Kellogg Observatory from 1937 to 1958. For many years he taught adult evening classes in astronomy at the museum.

Mr. Davis served on the three-man committee of experts that the American Astronomical Society chose in 1942 to establish a uniform pronunciation of star and constellation names. This led to an officially adopted list of approved pronunciations, which was reprinted in the March, 1958, issue of this magazine, page 220.

His versatility is indicated by the fact that he had practiced law for over 40 years, was a distinguished Arabic scholar, and had played professional baseball with the New York Yankees and the Boston Braves before World War I. At the time of his death, he was writing an extensive book on the history of constellation and star names.



One of the sections of the observatory's interferometer array, with a four-foot parabolic dish at the far left.

# NEWS NOTES

## TEMPERATURES ON MERCURY

The great difference in temperature between sunlit and permanently shadowed portions of the planet Mercury, which always turns one face toward the sun, has been investigated theoretically by J. C. G. Walker at Yale University. Two sources of heat are involved, solar radiation and the radioactivity of the planet's material.

On the sunward side, radioactive heating is very minor compared with the sun's direct rays. If Mercury radiated as a black body, the theoretical temperature at the subsolar point would be 621° Kelvin when the planet is at its average distance from the sun. During the 88-day orbital revolution, the computed subsolar temperature varies from 695° at perihelion to 565° at aphelion.

The night side is warmed primarily by internal radioactivity, with only a minor contribution from solar heat conducted through the interior. The radioactive heat generation has been evaluated by assuming Mercury to have the same composition as a chondritic meteorite, one gram of which liberates 1.6 ergs of thermal energy per year. Dr. Walker finds that the permanently shadowed surface is only at about 28° K. It was at one time suggested that meteoritic bombardment might supply enough energy to raise the night side's temperature appreciably, but the new calculations show this effect to be quite insignificant.

For Mercury's internal temperature distribution, the Yale physicist assumed that heat conductivity and the distribution of radioactive material are both uniform throughout the interior. The central temperature turns out to be nearly 5,000° K., slowly diminishing outward; it is about 2,000° at 300 miles below the surface. These figures are valid only for Dr. Walker's particular model, and are at best an approximation to conditions in the actual planet.


## NEW NATURAL SATELLITES OF THE EARTH?

The discovery of two faint, cloudlike objects circling the earth at the same distance as the moon has been reported by the Polish astronomer K. Kordylewski, at Krakow Observatory.

His find is the result of many years of searching, based upon the idea that the most likely locations for additional natural satellites are the Lagrangian points of the Earth-Moon system. These places have the property that small bodies located near them will persist in stable orbital motion. Two of these points, known as  $L_4$  and  $L_5$ , are at the same distance from the earth as the moon, the first 60 degrees ahead of the moon in its orbit, the other 60 degrees behind. In the Sun-Jupiter system, the corresponding

points are each the center of a group of Trojan asteroids.

Very few details are available concerning Dr. Kordylewski's discovery. His two libration clouds are both near the  $L_5$  point, several degrees apart, and were recorded on four photographs taken on March 6th and April 6th of this year. He suggests that similar objects may be found near the  $L_4$  point, 60° east of the moon. In *Circular 1760* of the International Astronomical Union, Dr. Kordylewski provides ephemerides for both libra-



As the moon travels around the earth, these  $L$ -points form equilateral triangles with the earth and moon. K. Kordylewski's two clouds are in the vicinity of the  $L_5$  point.

tion points, remarking that for observers in northerly latitudes the  $L_5$  clouds will be unfavorably placed until January, 1962, but the other point could be searched beginning this September.

As this issue goes to press, no observations of Dr. Kordylewski's clouds by other astronomers have been received.

## INTERFERENCE COATINGS ON ALUMINIZED MIRRORS

The reflection of visible light from metals can be increased by applying a pair of transparent dielectric films in the same manner as glass lens surfaces are coated to increase their transmission. In the Russian journal *Optics and Spectroscopy* (January, 1961), E. I. Levitina gives the results of experiments in which reflecting layers of aluminum, chromium, nickel, palladium, and rhodium were coated with films of silica and titanium dioxide.

In every case the over-all reflectivity was increased, from about 88 per cent to 91 per cent for aluminized surfaces such as commonly employed for telescope mirrors. But even greater advantage was gained in mechanical stability and resistance to deterioration. The Soviet scientist writes:

"After three to five hours testing in a corrosion chamber with periodic moisture

## IN THE CURRENT JOURNALS

N.A.S.A. LOOKS AHEAD, by Morton J. Stoller, *Spaceflight*, May, 1961. "Major Mercury flights will probably continue for several years. . . . Following the Atlas-Agena flights, the Saturn vehicle would be used for full-scale development and prototype flights. Earth-orbital missions, using the final spacecraft, could conceivably begin in 1966, with circumlunar missions following as soon as the state of both technical and aeromedical knowledge permits such flights."

THE TEMPERATURES OF THE PLANETS, by Cornell H. Mayer, *Scientific American*, May, 1961. "Various observers have now used radio techniques to measure the temperatures of Venus, Mars, Jupiter and Saturn. Some of the results are in line with advance expectations; others have come as quite a surprise."

condensation and a drop in temperature from 50° to 35° C. in the course of each hour (cycle), a layer of aluminum on glass completely disintegrates. After 100 hours an aluminum mirror of the same type, with films of silica and titanium dioxide, still does not show a trace of corrosion, and furthermore, the mirror retains the initial high reflection coefficient. The reflection coefficients of aluminum mirrors with films were lowered only to 89 per cent after exposure for one hour in a 10-per-cent solution of caustic soda, whereas a metallic layer on the mirror without films is completely dissolved under the same conditions in 30 to 45 seconds. Aluminum mirrors with films can remain for an hour in concentrated sulfuric acid without a change in reflection coefficient, whereas mirrors without films are completely dissolved in 20 to 30 minutes."

The deposited films of silica and titanium oxide were obtained by a relatively simple chemical process, the hydrolysis of ethyl esters of orthosilicic and orthotitanic acids.

## SHAPES OF LUNAR CRATERS

Undamaged lunar craters are to a first approximation circular, although the curvature of the moon's surface makes those situated near its limb appear quite elongated by the effect of foreshortening. But there is a systematic tendency for the craters to be slightly elliptical, according to Ernst J. Öpik, University of Maryland, partly as a result of changes in the rotation period and shape of the moon since they were formed. In the *Astronomical Journal* for March, 1961, he writes:

"According to the theory of tidal evolution, in the immediate past the moon was nearer to earth, when its tidal distortion may have been considerable. Impact craters formed at that time must subsequently have undergone deformation

when the moon assumed its present more nearly spherical shape.

"Actually, impact craters never are expected to be exactly circular; a slight ellipticity is predicted by the theory of impact, depending upon the angle of incidence and the 'explosiveness' of the collision. The [tidal-rotational] deformations superimposed on this random ellipticity will lead to elongation in a certain preferential direction, depending upon the location of the craters in relation to the tidal and rotational axes."

From measurements of 178 craters in G. P. Kuiper's *Photographic Lunar Atlas*, Dr. Öpik calculates that the observed crater ellipticities indicate an origin when the moon was between 30,000 and 50,000 kilometers from the center of the earth, roughly a tenth of its present distance. The period of revolution of the moon was much shorter then, and its rotation was faster, resulting in a more pronounced flattening at the poles. The tidal bulge was larger than at present. Hence a circular impact crater, formed on the moon at that time, would become deformed as our satellite receded, both because the polar flattening decreased as the rotation slowed down and because the tidal bulge

became less pronounced at the greater distance.

The Maryland astronomer believes that the larger lunar craters, considered in this study, probably were formed by an intense bombardment of short duration, possibly by earthbound fragments in a cloud or ring surrounding the geocentric orbit of the growing moon. Previously he had found (*SKY AND TELESCOPE*, March, 1960, page 285) that the number of craters less than 13 kilometers in diameter was consistent with meteoritic impacts continued over the past 4.5 billion years.

#### NAVAL OBSERVATORY LIBRARY

The largest library of any astronomical observatory in the United States is in Washington, D. C., at the U. S. Naval Observatory. It contains over 55,000 catalogued volumes, and many periodical sets. The collection was begun in 1811-15 by F. R. Hassler under the authorization of Thomas Jefferson, and thus is older than the observatory itself, which was established in 1830.

In charge of the Naval Observatory library is Marjorie S. Clopine, who writes in the February issue of *Special Libraries*: "The resources of the library make refer-

ence work especially rewarding. Typical reference questions concern such facts as: the official action ending the use of the astronomical day; coordinates of the observatory erected in Independence Square [Philadelphia] to observe the transit of Venus in 1769 (This observatory attained greater prominence as the structure from which the Declaration of Independence was first read publicly.); and literature about the surface of the moon, related to the problem of constructing an observatory thereon."

Besides being a very complete working assembly of modern astronomical literature, the Naval Observatory library also contains a notable collection of 500 rare astronomical works printed before 1800. Six of these are from the 15th century, the earliest being the first illustrated edition of Hyginus' *Poeticon Astronomicon* (1482), with its beautiful woodcuts representing the constellations.

Among other early books at the Naval Observatory is *Machina Coelestis*, by Johannes Hevelius, published at Danzig in two parts, 1673-79. The second part is extremely rare, for almost the entire edition was destroyed when a dismissed servant returned to set fire to Hevelius' house.

#### SOLAR MAGNETISM AND SUNSPOT FORMATION

(Continued from page 3)

self as a loop in which the lines of force exert a magnetic pressure comparable to the gas pressure. The magnetic buoyancy of the loop will make it rise and break through the solar photosphere, where it forms a bipolar magnetic region with a positive and a negative magnetic flux area.

The magnetic lines of force that link the two parts of the bipolar region loop up into the high solar atmosphere. Dr. Babcock's calculations show that ample magnetic rope is formed within the sun to account for the two or three thousand bipolar areas observed in each 11-year solar cycle.

Because of the opposite direction of twisting in the two hemispheres, the polarities of leading and following areas are interchanged, just as actually observed. The concentrated magnetic flux bundle piercing the solar surface inhibits local convection of the hot gases (as proposed by L. Biermann in 1941) and a sunspot may form. Sunspots are found only within bipolar magnetic regions, especially while the latter are young and compact. The observed spiral filaments in the chromosphere near spots may be explained by the twisting of the magnetic rope, and the duplicity of spots by the fact that the rope is looped up through the surface.


Expanding in the atmosphere, the leading part of a loop moves toward the equator as it ages, to be canceled by merging with vertical legs of other loops. Meanwhile the following part drifts

poleward, and in combination with other loops reduces and finally replaces (with opposite polarity) the original magnetic field of the sun. This process, which involves severing and reconnecting the lines of force in the corona, need be only one per cent efficient. It requires 11 years for completion, resulting in an unwrapped field of reversed polarity for the start of the next cycle.

As the vertical legs of the loops are canceled, they leave the upper parts drifting freely in the corona. These detached loops may travel outward through the

solar system carrying entrapped ions and electrons. By means of instruments carried by artificial satellites, such clouds of charged particles have been observed as they pass the earth.

Reporting his theory in the March issue of the *Astrophysical Journal*, Dr. Babcock characterizes the process as a freely running oscillator that lacks stabilization. The sun's differential rotation supplies only enough energy to drive it for a few thousand years. If this is indeed the driving force, the problem of the maintenance of that rotation assumes new interest.



Two stages in the solar cycle, according to Horace W. Babcock. At the left, the submerged magnetic lines of force have been drawn out in longitude and wrapped around the sun. At the right, magnetic flux loops have risen through the photosphere, each loop forming a bipolar magnetic region. Diagrams courtesy the "Astrophysical Journal."



# Radio Ursigrams

MARTIN H. POTTER

**M**ANY amateur astronomers engaged seriously in observing the sun lack current and accurate information from professional astronomers concerning solar happenings. Radio Ursigrams provide one method by which an amateur, with little expense or trouble, can obtain the latest reports on sunspots, prominence activity, auroras, ionospheric disturbances, and other related phenomena.

The name *Ursigram* comes from the French abbreviation for International Scientific Radio Union (U. R. S. I.). Ursigrams are messages, coded for brevity, that report data from 139 co-operating observatories and laboratories in all parts of the world. They are sent to users by radio, telegraph, and mail.

About 30 different types of Radio Ursigrams are distributed, each devoted to a specific class of information. Of these, CHRAG messages are especially valuable to amateur astronomers. They are transmitted daily from Ste. Assise, France, by station FYP, 91.15 kilocycles per second, at 12:08 and 13:08 Universal time. All broadcasts are in international Morse code (emission type A1), at a speed of

about 20 words per minute, and include other Ursigram messages besides the CHRAG bulletins.

A sample bulletin might read: CHRAG 17510 71221 78212. The decoded text tells: "Meudon Observatory, France, reports that on the 17th at 12 hours Universal time the sunspot number was 112, a high-quality determination. There was medium activity on the disk, and slight prominence activity. The sudden disappearance was noted of a prominence in the sun's western hemisphere, 80° of longitude from the central meridian, with a latitude between +20° and +30°. This same prominence had been observed one day earlier as of importance 2 on a scale of 1 to 3."


While a full key to the CHRAG code is too lengthy to be reprinted here, its nature will become clear from a partial explanation of this message. The first word, CHRAG, identifies the kind of message and the code used. The second group, 17510, tells first the day and tenth of day (Universal time) as 17.5, while 10 identifies the observatory as Meudon.

Information about the relative sun-

spot number is contained in the next group, 71221, which may be symbolized by WWWAP. If WWW in the message is from 000 to 299, it gives the sunspot number directly, but characterizes it as a poor determination. If WWW is between 300 and 599, the determination is of medium quality, and 300 must be subtracted to obtain the sunspot number; if between 600 and 999, the quality is high, and 600 must be subtracted. On the rare occasions when the actual sunspot number is greater than 299, WWW is still transmitted as 299, 599, or 999, depending on the quality of the observation. The true number is then given at the end of the message. The two remaining symbols A and P in this group are one-digit numbers characterizing, respectively, the level of general solar activity and the level of prominence and filament activity.

The fourth group, of five numbers, describes the location and characteristics of an activity center or prominence that has suddenly disappeared, and may be followed by other such groups.

Another series of Ursigrams uses the CHROM code to report solar chromospheric and photospheric activity. Others deal with the corona, solar radio noise, calcium plages, and further specific phenomena. Amateur users of the solar flare detectors described by David Warshaw (*Radio-Electronics*, January, 1959, page 41) or Philip Del Vecchio (*SKY AND TELESCOPE*, August, 1959, page 546) will find the UFLAR bulletins very profitable.



This photograph, showing prominences at the sun's limb typical of those reported in Ursigrams, was taken on June 26, 1956, with Walter Semerau's solar telescope in Kenmore, New York. The PROMO code would be used to report this activity in detail, giving prominence positions, importances, and times of occurrence.



## USSPA

### Sunspots

#### 1. — Content of the message

- Wolf relative number of sunspots.
- General activity indices for disk and limb.
- Area of spot groups.
- Position, importance and Brunner type of groups (Code A).

#### 2. — General form of the message

USSPA	JJTH	aaab	eeff	QXXYY	gijkk
(a)	(b)	(c)	(d)	(e)	(f)
key word					importance and number days group, Brunner classification and spot count
					quadrant, angular meridian distance and heliographic latitude
					serial number and area of spot group
					relative sunspot number, general activity indices for disk and for limb
					date and time of observation, identification

#### 3. — Definition of symbols

- (a) USSPA = key word (U = interchange, sunspot, Code A)
- (b) JJ = Greenwich date of observation  
T = time of observation in tenths of the Greenwich day  
II = indicator of the observatory (Table II)
- (c) aaa = relative sunspot number for date (observatories should give R + 600 if observation quality is good; R + 300 if fair, and R if poor)  
b = general activity index for disk, scale of 1 (small) to 5 (great) activity  
c = general activity index for limb 1-5
- (d) ee = serial number of spot group (assigned by reporting observatory on the basis of observations on preceding days)  
ff = area of spot group in tens of millionths of solar hemisphere
- (e) Q = quadrant (heliographic coordinates) containing the spot group (Table Q)  
XX = angular meridian distance, in degrees  
YY = heliographic latitude, in degrees
- (f) g = importance of group, scale 1, 2, 3  
i = number of days group observed since limb passage or first appearance on disk  
j = Brunner classification of region, A = 1, B = 2, etc.  
kk = spot count

#### 4. — Tables of Coding

##### (II) Indicator of the observatory

See Table (II), Section 2.

##### (Q) Quadrant

1 = NE	4 = NW
2 = SE	3 = SW

Note: Omit A, B, J type spots (URSI Information Bulletin, n° 94, Nov.-Dec. 1955).

An excerpt from the "Manual of Ursigram Codes," giving an explanation of USSPA messages. All groups after the code name are sent as numbers, for which the letters in section 2 are symbolic generalizations.

These permit quick evaluation of a flare's intensity and additional properties as observed at many other stations. Even if the user of these bulletins is unable to copy the code transmissions himself, he can probably find an amateur radio operator or experienced short-wave listener who is willing to help.


The schedules for the transmission of Ursigrams from data centers in France, West Germany, Japan, and India are published in the *List of Special Service*

*Stations* distributed by the International Telecommunication Union at Berne, Switzerland. Complete information regarding schedules, decoding methods for each of the 30-odd types of bulletin, and more, is contained in the *Manual of Ursigram Codes*, available from the General Secretary of U.R.S.I., 7 Place Émile Danco, Brussels 18, Belgium. The \$8.00 price for this loose-leaf book includes corrections and supplements, which are mailed to the user from time to time.

## QUESTIONS... FROM THE S+T MAILBAG

**Q.** In observing Mizar and Alcor with a small telescope, Mizar seems to split into two components. Is this star double?

**A.** Yes. In addition, both of the components, labeled A and B in the accompanying chart, are spectroscopic binaries. The star marked SL in the field diagram



is Sidus Ludovicianum, an 8th-magnitude star that can be seen in even small instruments.

**Q.** What is the limiting magnitude for visual and photographic observations with the 200-inch telescope?

**A.** Stars of about magnitude 19 can be seen with averted vision; the photographic limit is magnitude 23.

**Q.** How many telescopes are there in the United States having apertures of 20 inches or more?

**A.** According to G. P. Kuiper and Barbara M. Middlehurst's book *Telescopes*, there are 49 in this country. Of these, 33 are reflectors, 12 refractors, and four are Schmidt cameras.

**Q.** Was there a bright comet in late 1948 visible in the Southern Hemisphere?

**A.** Yes. This was Comet 1948I, which was first seen from South Africa during a total eclipse of the sun on November 1st, leading to the name "eclipse comet." It reached 2nd magnitude, and could be seen from the southern United States.


**Q.** How are rotation rates found for asteroids or satellites too small to show disks?

**A.** By measuring variations of a rhythmic nature in the light reflected by the body. If it is irregularly shaped, it will reflect different amounts of light as it turns, and the time required for these variations gives the period of rotation.

**Q.** How many natural satellites of planets in our solar system can be seen with a 10-inch reflector by a skilled observer under the most favorable conditions?

**A.** Possibly 13: in addition to our own moon, four of Jupiter, five of Saturn, two of Uranus, and one of Neptune.

W. E. S.



## E. E. Barnard and Milky Way Photography

OTTO STRUVE, *National Radio Astronomy Observatory\**

Edward Emerson Barnard (1857-1923) was one of the greatest astronomical observers of all time. Discoverer of 16 comets, he made numerous precise measurements of double stars, satellites, and star clusters. His most fundamental contribution was to employ photography to study the Milky Way. Yerkes Observatory photograph.

no other astronomers observed it. Hence the first comet officially bearing his name is the one he detected on September 17th that year in Virgo. He continued for many years searching the sky, eventually discovering a total of 16 comets. On five different occasions, Barnard received the \$200 prize given by H. H. Warner, of Rochester, New York, for each unexpected comet discovered by an American observer. This made it possible for Barnard to buy a modest home for himself and his family, which was known in Nashville as the "comet house."

In 1883 Barnard was appointed to a small fellowship at Vanderbilt University in Nashville, which enabled him to attend courses while taking care of its observatory. He received the bachelor of science degree from this university in 1887.

That same year, E. S. Holden, the first director of Lick Observatory, offered Barnard a position at the new institution, where he remained until 1895. One of his greatest discoveries with the 36-inch refractor was the fifth satellite of Jupiter. This faint moon was found on September 9, 1892, and observed by Barnard on many occasions both at Mount Hamilton and Yerkes Observatory. Even with very large telescopes it is an extremely difficult object, visible only when seeing and transparency are excellent.

Barnard's most important contribution was his astronomical photography with varied types of instruments, especially wide-field photographic objectives. He revolutionized the art, although experiments in the application of photography to astronomy had started before Barnard was born.

The daguerreotype process was invented in 1839, and within a year the American amateur J. W. Draper had succeeded in obtaining a picture of the moon. Because the daguerreotype method and the wet-plate process (invented about 1850) were very slow, it seemed at first that the only hope for any success lay in photography of the sun, moon, and possibly the planets and some of the brighter stars. There was no prospect of photographing faint stars or nebulae.

One difficulty in the early attempts was

WHEN I ARRIVED at Yerkes Observatory in 1921 as a refugee from war-torn Russia, one of the first persons I met was an elderly gentleman who was looking at the barograph in the observatory library, and quietly complaining that the barometric pressure was going down and the night would presumably be cloudy. This was Edward Emerson Barnard, the most famous member of the Yerkes Observatory staff, generally recognized as the most capable and productive astronomical observer in the world. I knew him for only 18 months, for he died on February 6, 1923, in his 66th year.

Although he published some 900 scientific papers, his name is now seldom mentioned in astronomical literature. Nevertheless, all astronomers active today are in one way or another building upon the foundation of his accurate measurements of celestial bodies, and his thousands of Milky Way photographs, obtained at Lick Observatory and later at Yerkes. It is surprising that references to Barnard's contributions are not more often cited by those who have followed in his footsteps. His name and work have become part of astronomical history sooner than is normal at the present time. This has happened, perhaps, mainly because later generations of astronomers found nothing to criticize in Barnard's work, and associated him with the great names of earlier decades.

Barnard was born in Nashville, Tennessee, on December 16, 1857, and throughout his life he remembered the flashes of artillery at the battle of Nashville during the Civil War. In his youth

he survived an attack of cholera when that plague raged across the southern states.

Because of the need to support his impoverished family, he attended public school for only two months, beginning work as a photographer's assistant in Nashville at the age of nine. His principal duty was to point a heavy enlarging camera at the sun. To make the work easier and to save time for reading and study, he built himself a simple equatorial mounting with a driving mechanism, so that the camera automatically followed the sun's diurnal motion. By 1877 he had saved enough money to purchase a 5-inch equatorial refractor, and he was already well-known as an active amateur astronomer when shortly afterward he joined the American Association for the Advancement of Science.

At the 1877 meeting of the association in Nashville, Barnard met the famous Simon Newcomb, director of the U. S. Nautical Almanac Office, and at the age of 42 a leader in mathematical astronomy. Newcomb advised Barnard to secure a firm grounding in mathematics, but the latter's limited resources restricted him to studying at home. While Barnard became proficient in elementary mathematics, he was never thoroughly acquainted with the more advanced branches, and his fame rests upon his success as an observer rather than as a theoretician.


On May 12, 1881, at the age of 23, Barnard discovered his first comet in the morning sky near Alpha Pegasi, but the object was not formally announced, and

\*Operated by the Associated Universities, Inc., under contract with the National Science Foundation.



The intricate splendor of the Milky Way was first revealed by Barnard's long-exposure photographs. He took this picture of the Sagittarius starclouds with the 6-inch Willard lens at Lick Observatory on August 13, 1895, a three-hour exposure. Barnard called these starclouds the finest in the sky, so full of splendid details as to defy adequate description. This view is centered at right ascension  $17^h 55^m$ , declination  $-28^\circ$ , with north toward the top and a scale of  $1\frac{1}{2}$  degrees per inch. The very bright star in the lower left corner is Epsilon Sagittarii, while Delta (dim because of its redness) is one inch from the left edge and five inches from the top. The star clusters M6 and M7 are at lower right. The nebulous cluster M8 is conspicuous  $1\frac{1}{2}$  inches from the top edge and  $3\frac{1}{4}$  inches from the left, with the Trifid nebula, M20, above it and slightly to the right. The cluster M21 is above the latter at the very top of the picture. Seen in silhouette against the bright starry background are many dark wisps and patches. One of these, looking like a tiny ink drop, is just above the picture's center. Three inches below it is another small dark nebula, likened by Barnard to a parrot's head, with a star for the eye. Reproduced from Lick Observatory "Publications," Vol. XI (1913).





The Bruce instrument at Yerkes Observatory as it appeared when Barnard used it. Three telescopes are rigidly mounted together. Seen fully in the center is a 6½-inch f/5 Voigtländer, and above it on the left a 5-inch guiding telescope. Behind the latter, mostly hidden, is the main 10-inch Bruce camera, whose 12-inch plates cover a field 13½ degrees square. A 3.4-inch Clark lens of 20-inch focus is in the auxiliary camera at upper left.


that existing refractors were corrected for the yellow part of the spectrum to which the eye is most sensitive, whereas the photographic emulsions were sensitive to blue light. L. M. Rutherford in 1864 made a great step forward by making the first telescope lens corrected especially for photographic wave lengths. Another major advance was the introduction of the relatively fast dry plate about 1880. However, in 1900 Barnard wrote, "What is shown by the best lunar photographs has not yet approached that which can be seen with a good telescope of very moderate size." In a certain sense this remains true even today.

The first daguerreotype of the sun and its spots was obtained in 1845 by H. L. Fizeau and L. Foucault in France, and Berkowski at Königsberg Observatory took the first picture of the solar corona during the total eclipse of July 28, 1851. By 1900, photography was used extensively in solar work, and many photographs of the planets, comets, and asteroids had been obtained. The greatest advances had been made, however, in the field that

at first seemed to hold no promise for photography — the stars and nebulae. This had come almost as a surprise to astronomers.

In the fall of 1882, a bright comet ap-

Compare this picture of a group of dark clouds in the Aquila Milky Way with three other views of the same field on the facing page. F. E. Ross used a 5-inch f/7 lens of his own design for this four-hour exposure on July 17, 1931, at Mount Wilson Observatory. The faintest stars in this reproduction are about 16th magnitude. From the Ross-Calvert "Atlas of the Northern Milky Way," University of Chicago Press (1934).




peared in the southern sky, inducing David Gill at the Cape of Good Hope to borrow a camera from a local photographer. His pictures showed the comet well, but even more impressive was the great number of stars recorded on the negatives. At this time, the Paris astronomers Paul and Prosper Henry, who were making maps of faint stars along the ecliptic, had reached the Milky Way and were discouraged by the immense numbers of stars encountered. On seeing Dr. Gill's pictures, they realized the potentialities of photography for making charts, and it may be said that the active application of stellar photography began with this incident.

As for nebulae, the first photograph of the most conspicuous one visually, Messier 42 in Orion, was made in 1880 by Henry Draper, son of J. W. Draper. Before the Henry brothers had constructed their first photographic telescope, much controversy had raged over the existence of a nebula connected with the star Merope in the Pleiades, which was promptly confirmed by their photographs. According to Barnard, one of these "showed the Merope nebula just as the best observers had drawn it and at the same time filled the entire group of stars with an entangling system of nebulous matter which seemed to bind together the different stars of the group with misty wreaths and streams of filmy light, clearly all of which is beyond the keenest vision and the most powerful telescope. This was a revelation. . . . It began to dawn upon astronomers what great possibilities lay in the photographic plate for the detection and study of nebulae."

In the 1880's, the best photographs of nebulae were obtained by A. A. Common in England, with his 36-inch reflector. Barnard has stated that a nebular photograph by Common and one of a rich star field in Cygnus by the Henry brothers first called his attention to the astronomical value of photography, and that thus he conceived the idea for his proj-





Two more views of the dark clouds northwest of Altair, whose overexposed image is near the lower left corner of the first picture. This was taken by Barnard on October 12, 1892, an exposure of four hours 20 minutes with the 6-inch Willard lens. Comet 1892 V was discovered on this photograph; its faint, fuzzy trail is nearly an inch from the top,  $1\frac{1}{2}$  inches from the right edge. The second photograph is also by Barnard, a four-hour exposure with the 10-inch Bruce telescope on August 27, 1905. The C-shaped dark marking just below center is No. 143 in Barnard's catalogue of obscuring nebulosities. Slightly to the right and below is his No. 142. To the left of the former, just at the edge of the picture, is another, less conspicuous dark nebula, Barnard 340. The peculiar shapes of many of these markings led him to the correct conclusion that they are opaque foreground objects, rather than vacancies in the star clouds.

ect of photographing the Milky Way.

He soon realized the advantages of a fast, wide-angle camera over ordinary telescopes for recording the Milky Way. His early work at Lick was done with the 6-inch Willard lens (named after a photographic stock dealer), which dated back to the days of wet-plate portraits, when a camera of great light-gathering power was needed to shorten the time of portrait sittings. Lick Observatory purchased this lens with funds donated by C. F. Crocker, after it had been used by an amateur astronomer to take excellent photographs of the solar eclipse of January 1, 1889. Volume XI of the Lick Observatory *Publications* (1913) contains 129 photographs, mainly of the Milky Way and comets, made by Barnard with the Willard lens. In addition, many of his fine pictures appeared in various issues of the *Astrophysical Journal*.

Barnard's last series of photographs of our galaxy was published in 1927 as *A Photographic Atlas of Selected Regions of the Milky Way*. This included 50 long exposures obtained with the 10-inch Bruce refractor at Yerkes and at Mount Wilson observatories. (He had spent part of 1905 as a guest investigator at Mount Wilson, taking the Bruce telescope with him.) This instrument and its dome were erected with money donated by Catherine W. Bruce, a generous patron of astronomy who had provided several observatories with large photographic telescopes. It

consists of two f/5 cameras of 10 and  $6\frac{1}{4}$  inches aperture, rigidly bound together on the same mounting with a 5-inch guiding telescope.

To maintain the highest possible quality of the atlas, Barnard himself made the reproducing negatives, and personally inspected each of the more than 35,000



Part of the same field in Aquila, photographed with the 48-inch Schmidt telescope of Palomar Observatory on August 12-13, 1958. Hold the picture at arm's length to accentuate the dark markings, especially the delicate lanes left of Barnard 143. Mount Wilson and Palomar Observatories photograph.



The Milky Way from Aquila to Carina, photographed by A. D. Code and T. E. Houck in infrared light with a Greenstein-Henney wide-angle camera. Note the distinct belt of obscuration, much as in an edge-on view of a spiral galaxy. The three radial shadows are caused by the three-legged camera support. Reproduced from the "Astrophysical Journal," March, 1955.

prints required for an edition of 700 copies. He had drafted detailed descriptions of the individual pictures before his death in 1923, and the work was completed for publication by his personal assistant and niece, Mary Calvert.

An amusing incident occurred during the preparation of the atlas prints. They were all made in Chicago and shipped to Yerkes Observatory by truck. During one of the Chicago gangster wars, a shipment of photographs was completely pierced by a bullet. Of course, the pictures had to be replaced, but the damaged ones for a long time remained objects of curiosity to visitors at Yerkes Observatory.

An important feature of the *Atlas* was the key charts that identified all the brightest stars and the dark and luminous nebulae. Barnard included in the introduction a catalogue of 349 dark nebulae. Awareness of these dark regions in the Milky Way can perhaps be traced back to Sir William Herschel's expression, "Surely, this is a hole in the heavens!" But their first systematic study and correct interpretation are due to Barnard.

In 1919 he wrote: "I did not at first believe in these dark obscuring masses. The proof was not conclusive. The increase of evidence, however, from my own photographs convinced me later, especially after investigating some of them visually, that many of these markings were not simply due to an actual want of stars, but were really obscuring bodies nearer to us than the distant stars. In this way it has fallen to my lot to prove this fact. I think that there is sufficient proof now to make this certain."

Barnard's catalogue of dark clouds was

the only one until 1960, when an atlas of galactic dark nebulosities by D. S. Khavtasi was published in the Soviet Union. This consists of six charts containing stars down to apparent magnitude 6, and with a co-ordinate grid showing galactic latitude and longitude. Apparently Khavtasi used the Ross-Calvert atlas and southern Milky Way photographs. The dark nebulae are shown by shaded areas, indicating three different amounts of opacity, and are labeled with arbitrary numbers ranging from 1 to 762. However, many of the numbered objects are really condensations in larger obscuring aggregates.

As has long been known, the dark nebulae tend to lie close to the central line of the Milky Way. In the Soviet atlas, the widest distribution in galactic latitude occurs in the vicinity of the galactic center, where the dark features are spread approximately uniformly between latitudes  $-10^\circ$  and  $+15^\circ$ . In the region of Cygnus and Vulpecula, this range is from about  $-7^\circ$  to  $+5^\circ$ .

G. A. Shajn and V. Hase had found in 1952 that the dark clouds of the Milky Way are generally elongated parallel to the galactic plane. (Usually this has been interpreted in terms of an interstellar magnetic field whose lines of force lie predominantly in the galactic plane.) Such a trend is also noticeable in the new atlas, but individual clouds often depart from it strikingly.

Another catalogue of dark nebulae is now being prepared by Mrs. B. T. Lynds from the Palomar sky survey photographs taken with the 48-inch Schmidt telescope.

All of Barnard's Milky Way photography was carried out with lenses rather than mirrors. Bernhard Schmidt did not

invent the wide-angle reflecting telescope until 1932, after Barnard's death. At the present time, the best photographs of large Milky Way regions are being obtained with either Schmidt or Maksutov-type reflectors. The Palomar Schmidt telescope has a 48-inch correcting plate and a spherical mirror 72 inches in diameter. A 32-inch Schmidt instrument is at Bergedorf Observatory, near Hamburg, West Germany. Several large Maksutov telescopes are operating in the Soviet Union, for example at the Crimean Astrophysical Observatory and at Burakan Observatory in Armenia, and excellent photographs have been taken with them. Recent experiments seem to indicate that the optical performance of the Schmidt telescope is somewhat better than that of the Maksutov, but both types are excellent for covering sky fields of the order of 20 degrees across. Small wide-angle optical systems developed by L. G. Henney and J. L. Greenstein have recently proven very useful. Shown here is a typical Milky Way photograph taken with a Henney-Greenstein camera, covering a region of the sky 140 degrees across.

Barnard was primarily interested in the shapes of the dark cosmic clouds. Max Wolf of Heidelberg, who also contributed significantly to astronomical photography, invented a statistical method to obtain crude estimates of the distances and opacities of different dark nebulae. Later investigators found that starlight passing through obscuring clouds is appreciably reddened (which indicates that the interstellar particles in the clouds have sizes of the order of  $1/100,000$  centimeter) and that the transmitted starlight is often partly polarized.

Stars happening to lie near a dark cosmic cloud may illuminate parts of it, thereby producing reflection nebulae whose spectra resemble the stars'. A reflection nebula lit by a B-type star is strongly blue, whereas the dense cosmic cloud near Antares is almost as red as Antares itself. The nebula surrounding the Pleiades stars was the first one to be recognized as of the reflection type, by V. M. Slipher at Lowell Observatory in 1912.

Although photographs of dark nebulae have been accumulating for 70 years, we still have no reliable information concerning the motions of these objects. But there is every reason to believe that we shall soon be able to detect displacements of the edges of certain dark clouds, and probably also the gradual fading of stars passing behind them, or the emergence of previously invisible objects. Long ago, attempts were made to explain certain types of stellar variability by such processes, but there is yet no definitely established case of a star whose brightness has been altered by changes in the opacity of the foreground material.

I am grateful to Mrs. V. Zebergs for her assistance in preparing this article.


# GETTING ACQUAINTED WITH ASTRONOMY

## JUPITER'S SATELLITES

WITH a dozen known satellites, Jupiter is the center of a miniature solar system. No other planet has so numerous a retinue of secondaries (if we omit the tiny artificial bodies circling the earth and the host of moonlets that make up the Saturnian ring system). The runners-up are Saturn with its nine moons, and Uranus with five.

Jupiter's satellites are of two sharply contrasting kinds. Four of them are giant moons, comparable in size to the earth's satellite or to the planet Mercury, and are bright enough to be seen readily in binoculars. They were among the first objects discovered with the newly invented telescope in the winter of 1609-10, independently by Galileo in Italy and by Simon Marius in Germany. Recent historians have established that Marius was the first to note their existence, and the fact of their revolution around Jupiter, but Galileo had priority in recognizing the fourth satellite. Together, these bodies are often referred to as the Galilean satellites; individually, they are designated I, II, III, and IV, in the order of increasing distance from the planet. The corresponding names, Io, Europa, Ganymede, and Callisto, were introduced by Marius.

The remaining eight satellites are small and very faint. With the 36-inch Lick refractor in 1892, E. E. Barnard discovered V, a little point of light even closer to Jupiter than I, orbiting in only 11 hours 57 minutes. Because of the overwhelming glare of the planet, this difficult object is observable only in very large telescopes. Satellites VI to XII are faint moons moving far outside the others, and were discovered photographically between 1904 and 1951. Their magnitudes range from




Ganymede, Jupiter's third satellite, as seen in the 15-inch refractor of the Jarry-Desloges Observatory, on six nights between December 12, 1917, and February 13, 1918. Changing marks indicate that the moon rotates.

14 to 18. Remarkably, four of them (VIII, IX, XI, and XII) are moving around Jupiter in the opposite direction from all the others. The outer satellites are so remote from the planet that their motions are subject to large disturbances from the sun's attraction, posing difficult problems for specialists in celestial mechanics.

The four Galilean satellites, on the other hand, present a variety of interesting phenomena for amateur observers. Because of their relative proximity to Jupiter, I, II, and III on every revolution pass in front of the planet's disk, behind it, and through the Jovian shadow cone. In addition, the shadows of the satellites themselves move across the disk. But IV, in its larger orbit, shows these phenomena only during about five years out of every 12, at other times passing north or south of the disk and shadow cone.

For a specific illustration, consider the events of I during the first days of July. This is before the planet's opposition (July 25th); hence, the shadow cone of Jupiter extends to the west of its disk in the sky, as the chart on page 55 indicates. On July 1st at 5 o'clock Universal time, I is in the farther half of its orbit, slightly west of the disk and approaching it. At 5:20, the moon disappears into eclipse. Because of the appreciable size of the satellite, it takes about two minutes to fade from bright 6th magnitude to invisibility — a striking sight!

Io's exit from the shadow will be unobservable, taking place behind Jupiter. This moon next becomes visible at 8:13, as it emerges from occultation at the planet's east limb. In the United States, all of the foregoing events occur during the hours of darkness on the night of June 30-July 1, the correction from Universal time to standard time being five




Jupiter and its four Galilean satellites, as they appear in a small telescope. From "Astronomie," by L. Rudaux and G. de Vaucouleurs.

hours for East Coast observers and eight hours for those on the West Coast.

During the following hours, I moves out to eastern elongation from Jupiter, then approaches the disk again while traversing the nearer half of its orbit. On the 2nd at 2:28 UT, the shadow of I enters upon the eastern edge of the disk, followed by the satellite itself 34 minutes later. In a telescope, the shadow appears as a very small, very black dot, and the moon as a tiny light disk. They travel westward together across Jupiter's face, leaving it at 4:46 and 5:20, respectively. Thereafter, Io travels out to western elongation, then swings back to enter eclipse again at 23:49 UT, July 2nd. The whole cycle has taken 1 day 18 hours 29 minutes — the synodic period of I.

The pattern just described is typical of I and II before Jupiter has reached opposition. After opposition, the planet's shadow is displaced eastward of the disk. This changes the sequence of events to the following: beginning of occultation at the west limb, reappearance from eclipse east of the planet, eastern elongation, beginning of satellite transit, beginning of shadow transit, end of transit, and



This model of the orbits of Jupiter's 12 satellites, made by Owen Gingerich, is viewed from a point above the planet's pole. The paths of the inner moons are drawn on a sheet of plastic, and those of the outer satellites are of brass wire supported by dowels. The actual orbits of the outer satellites change considerably from one revolution to the next, because of large perturbations by the sun.




end of shadow transit. At the time of opposition, Jupiter's shadow is concealed behind the disk, so for a few days eclipses will be practically unobservable.

An additional possibility arises for Ganymede and Callisto when Jupiter is several months from opposition, and its shadow cone has a considerable eastward or westward shift. Under these circumstances, both the beginning and end of an eclipse or occultation can be observed. For example, in August this year, IV undergoes occultation from 21:00 on the 24th to 1:28 on the 25th, and is visible on the east side of the planet until eclipse from 3:45 to 8:18.

Predictions of these events are published yearly in the *American Ephemeris* (on pages 344 to 365 of the volume for 1961). The predicted times are only rough approximations, and sometimes differ by several minutes from observation. Certain rare events, such as the eclipse of one satellite by another, are not listed there, but are forecast in the *Handbook* of the British Astronomical Association (see page 32 of this issue).

Easiest of all satellite phenomena to



Some of the phenomena presented by Jupiter's satellites are shown in this diagram for a time when Jupiter is near western quadrature to the sun. From the earth, III would be seen passing in front of the planet (in transit). Both I and its shadow, indicated by the dashed line, are also in transit. II is occulted by Jupiter, while IV is eclipsed by the planet's shadow.

observe are the eclipses, for which a 3-inch refractor serves well. Timings of these events can be recorded to 0.1 minute, noting the last speck of light at a disappearance, or the first speck at a reappearance. Because of the slower motions of III and especially IV, their light changes are more gradual, making first or last visibility more uncertain than for I or II. Since the threshold of visibility depends on many factors, eclipse timings must always include a record of the instrument used and sky conditions.

Transits of satellites across the disk of Jupiter require, as a rule, larger instruments for satisfactory viewing. Since the edge of Jupiter is appreciably darker than its center, a satellite usually appears lighter than the planetary background just after entry, and may look darker than the background during mid-transit. High powers and good seeing are desirable for watching shadow transits. When the shadow is just entering or leaving the disk, it appears greatly distorted against the curved surface of the planet.

In 1961, the orbits of the Galilean satellites are presented nearly edgewise to the earth, and a transiting or occulted satellite moves along the planet's equator. This makes possible a fairly accurate evaluation of the equatorial diameter of Jupiter by timing the duration of an occultation or transit. Suppose a transit of I is found to last 138 minutes, or 0.0541 of the synodic period of 2,549 minutes. Multiplying this fraction by the circumference of the orbit, 1,648,000 miles, gives 89,200 miles for Jupiter's diameter.

For the convenience of amateurs wishing to try this method, the circumferences of the other satellite orbits are: II, 2,621,000 miles; III, 4,181,000; and IV, 7,355,000. The corresponding synodic periods are: II, 5,118 minutes; III, 10,320; IV, 24,125.

If enough optical power is employed to show the satellite's disk clearly, it is possible to time the interval between external and internal tangency of its disk and Jupiter's. Then the same kind of calculation can be made for the diameter of the satellite.

The angular diameters of the Galilean satellites are, according to E. E. Barnard's micrometer measurements with the Yerkes 40-inch telescope: I, 1.05 second of arc; II, 0.85; III, 1.51; and IV, 1.43. The disk of the largest moon, Ganymede, can be made out with a 4-inch telescope under very favorable conditions. R. T. A. Innes, who made regular observations of these satellites from 1908 to 1925 with a 9-inch refractor in South Africa, reported that 700x would always show him the disks, but that this was not the case with 400x. On many occasions, a satellite entering eclipse appeared dichotomized, that is, like a tiny quarter moon.

Surface features on the satellite disks can be usefully observed only with very large apertures. Barnard with the 36-inch Lick refractor at 1,000x found that I has a bright equatorial belt and dark polar caps, while III and IV have bright polar caps and conspicuous though vaguely shaped dusky markings. The easiest of these features is III's bright cap, which has been definitely seen with a 12-inch telescope.

The rotation of these mottled disks produces small brightness variations which, according to J. Stebbins' photoelectric measurements in 1926-27, amount to 0.21, 0.30, 0.16, and 0.10 magnitude, for satellites I to IV, respectively. In each case, the period of light change is equal to that of orbital revolution. Thus each of these bodies always keeps the same face turned toward Jupiter, in the same way that our moon behaves relative to its primary, the earth.



Satellites X and XI, photographed with the 100-inch telescope by Seth B. Nicholson. The lower pair is from the discovery plates of XI, taken on July 30, 1938, exposure 60 minutes. The satellite's motion is shown by its change in position among the stars in the field. Mount Wilson and Palomar Observatories photographs.



## LETTERS

Sir:

During the February elongation of Mercury, I saw that planet with the naked eye on seven consecutive dates, from the 6th to the 12th. Altogether, this makes 57 days in the last 10 years on which I have seen Mercury with the naked eye or a telescope.

This record attests to the transparency of the New Mexico atmosphere. All observations have been made at Alamogordo or the neighboring Air Force Missile Development Center. On any good night stars are visible to magnitude 6.2, and to 6.4 when conditions are exceptional. I have observed Uranus with the naked eye on four evenings.

Venus has been seen 11 times by daylight, the latest sightings being on February 19th, 21st, and 23rd. Most of the navigational stars and bright planets can be viewed by day with standard 20x theodolites of 1½-inch aperture.

BERNARD D. GILDENBERG  
901 Cauthen Lane  
Alamogordo, N. M.

Sir:

Recently, Otto Struve described some of my work on radio source Cygnus A for the readers of *SKY AND TELESCOPE* (November, 1960, page 259), and in a subsequent letter R. Minkowski described his views on the subject (January, 1961, page 23). My paper, "Radio Galaxies," was circulated in the Soviet Union in the middle of January, and an English translation of it is being published in the current issue of *Soviet Astronomy — AJ*.

In it I emphasize that Cygnus A is most likely a double galaxy with the nuclei forming a very close pair, and that the components are genetically related. As another, but much less likely possibility, I consider the hypothesis that Cygnus A represents a single galaxy crossed by a broad dark band, similar to NGC 5128, which is identified with Centaurus A. Although I advanced this hypothesis as early as 1953, I had myself long since felt it to be improbable, and Dr. Minkowski has argued strongly against it.

Cygnus A and Centaurus A are similar, I believe, not merely in the resemblance between their associated optical objects, but in having analogous radio structures. Each has two radio-emitting regions, located symmetrically on opposite sides of the galaxies observed optically. These radio-emitting clouds are highly unstable formations, for there is no force in nature that can prevent them from expanding indefinitely. Their surface intensities are, therefore, diminishing.

Extrapolating backward in time for Centaurus A, we find that its radio luminosity was once thousands of times as great as now, and its brightness tens of thousands of times. Some hundreds of millions of years ago, it was a "twin" to

Cygnus A as the latter radio source appears now.

It has recently been found that the extended source of small angular size associated with the dark band in NGC 5128 is double, with the line joining the two centers lying within the band. Possibly this effect of doubleness arises from an edge-on view, with an absence of radio emission at the center of the system. The latter situation could be due to a high density of interstellar gas, which would rapidly destroy relativistic particles that produce radio emission by the synchrotron mechanism. A similar phenomenon may perhaps occur in the nucleus of our own galaxy.

In my paper, I advance a number of arguments against the colliding-galaxy idea, none of which Dr. Minkowski has effectively refuted. A number of astronomers now working in this field either categorically reject the collision hypothesis (V. A. Ambartsumian opposed it even earlier than I) or consider that it is by no means the only permissible one. Instead, the nature of radio galaxies seems intimately connected with the origin and evolution of optical galaxies.

In the early "halo" stage of the formation of a galaxy, Type I supernovae must occur frequently, an immense number of relativistic particles being formed in a relatively short time as by-products of the nucleogenetic process. Objects such as Centaurus A and, especially, Cygnus A must be relatively young. Yet most radio galaxies appear to be stable objects. This means that the relativistic particles occupy a rather considerable volume around each such galaxy and do not dissipate indefinitely out into extragalactic space.

The relativistic electrons are here formed through collisions of very fast heavy particles with interstellar gas nuclei. Since these processes occur quite slowly, the intense radio emission of a young galaxy may persist over billions of years. In this event, the radio power is not determined by the "instantaneous" frequency of supernova outbursts in a given galaxy, but by their frequency early in its history.

We still do not know whether the formation of jets — radio-emitting clouds consisting of relativistic particles, magnetic fields, and gas — is a peculiarity of such powerful objects as Cygnus A, Centaurus A, and a few others, or whether it occurs in practically every young galaxy. At any rate, some of the very fast particles produced during nucleogenesis, perhaps even the majority, are not released to extragalactic space.

Furthermore, what has become of the relativistic particles formed during nucleogenesis in our own galaxy? There are reasons to suppose that they were once hundreds of times as abundant as now. Most likely they have been destroyed in nuclear collisions; our stellar system is

quite old, and there has been enough time for this. A state of equilibrium now prevails, with the Milky Way's radio power being determined by the "instantaneous" frequency and power of supernovae. Perhaps this condition indicates that our galaxy is normal with regard to its radio power.

The foregoing illustrates my belief that the problem of the radio galaxies is closely interwoven with those of cosmogony and cosmology. The radio-emitting power of any galaxy seems to be governed by fundamental laws inherent in its evolution.

I. S. SHKLOVSKY  
Sternberg Astronomical Institute  
Moscow, U. S. S. R.

ED. NOTE: The foregoing is a condensed translation of Dr. Shklovsky's letter, which was written in Russian. The pertinent thoughts of the original have been preserved, but some of the technical details of the discussion have been omitted as they are more subject to direct correspondence among the scientists concerned.

Sir:

I have come upon a vivid account of a "wonderful display" of the aurora borealis, seen from near Rochester, New York, on September 22 and 23, 1827. Can some reader furnish me with other reports of this occurrence?

RUTH SHINSEL  
1610 M Ave.  
La Grande, Ore.

Sir:

The Ceylon Astronomical Association was founded in June, 1959, and we now have 56 active members. Our journal, *Equatorial*, first appeared last October. Its name is based on the fact that Colombo is only seven degrees north of the equator.

Under the direction of E. E. Vezey, formerly of Texas Agricultural and Mechanical Arts College, several members of our society have been grinding mirrors at home. I recently completed a 10-inch Newtonian reflector, and two others finished 8½-inch instruments. The chief difficulty in telescope making in this country is obtaining glass of the necessary thickness.

Ceylon has several small observatories; two of them own mirrors by the former director of Armagh Observatory, Reverend William F. A. Ellison. A 12½-inch mirror made by him is in the observatory at Kandy, the historic hill capital of Ceylon, and Colombo possesses a 13-inch reflector.

Since we are quite new to astronomy, we would like to start corresponding with more experienced amateurs and groups around the world, in order to exchange ideas and techniques.

D. S. HERSCHEL GUNAWARDENA  
31/5 Castle St.  
Colombo, Ceylon

# Johann Kepler and the Laws of Planetary Motion

RUFUS SUTER

**I**F YOU stick two tacks through a sheet of paper, tie a thread loosely around them, and, with a pencil holding the thread loop taut, trace the resulting curve, you will draw an ellipse with the tacks as its foci.

Kepler's first law of planetary motion states that a planet travels in an ellipse with the sun at one of the foci. The publication of this law in 1609 marked a very important advance in astronomy. All of Kepler's predecessors — whether like Ptolemy they believed the heavenly bodies to move around the earth, or like Copernicus to move around the sun — held that celestial bodies swung in perfect circles. Even Galileo seems to have regarded Kepler's ellipses as visionary nonsense. But Kepler found from Tycho Brahe's observed positions of Mars that its orbit was in fact an ellipse, and this led him to state his first law.

Another ancient prejudice, that each planet moves at an unvarying speed in its orbit, was refuted by Kepler's second law, also announced in 1609. According to this principle, the radius vector of a planet (the line joining it to the sun) sweeps over equal areas in equal intervals of time. In the diagram, the shaded areas are all equal; hence the planet requires the same length of time to move from A to B in its orbit as it does from C to D, or from E to F. Thus it travels fastest when nearest the sun (at perihelion), and slowest when farthest from the sun (aphelion).

In this second law, Kepler made a major innovation. Previous astronomers had treated the apparent motions of the heavenly bodies as a problem of geometric perspective. Kepler's approach was in a sense that of an engineer rather than a geometer. While he was interested in the purely geometric aspects of planetary motions, he also pondered the mechanical causes of these motions. Consideration of



Johann Kepler (1571-1630) inherited the observations of Tycho Brahe, and from them derived his laws of planetary motion. Kepler's search for cosmological patterns inspired Paul Hindemith's recent opera, "Die Harmonie der Welt."

how the planets should move if there were a center of power in the sun gave him his first inkling that each planet travels faster when nearer the cause of its motion. In the older approach, it was possible for an astronomer to deny that he asserted anything about the *real* movements of the heavenly bodies when he adopted the Copernican or heliocentric hypothesis. He was merely choosing one of several sets of geometrical patterns for simplicity in calculation or for subjective aesthetic reasons. This expedient became quite popular among scientists when the issue of the heliocentric theory versus theological authority arose. Kepler, with his dynamical point of view, could no more use it than could an engineer in a

later age believe that a wheel connected to a steam engine by a belt made the engine run, rather than vice versa.


Kepler's third law, announced in 1619, is a relation between the revolution times of the planets and the sizes of their orbits. It states that the ratio of the squares of the periods of any two planets is the same as the ratio of the cubes of their mean distances from the sun. In other words, with the earth's year and distance from the sun as units, the period squared for any planet is equal to its distance cubed. Thus, one moving in a circular orbit at four times the earth's distance from the sun would have a period equal to the square root of  $4^3$ , or eight years.

These three laws made possible much better predictions of planetary positions and eclipses of the sun and moon. They could be used for drawing up accurate ephemerides, and for fixing dates in ancient history. Later in the 17th century, Newton's law of gravitation, from which Kepler's laws are deducible, gave a rational and unifying explanation of them. It could then be seen that Kepler's laws are actually close approximations, which would be exact if the masses of the planets were negligible compared to the sun's.

Some of Kepler's contemporaries, among them Galileo, were seeking a physical proof for the theories of the earth's motion and the central position of the sun. Few at that time recognized it in the Keplerian laws. Nevertheless, from our hindsight we can see today that the proof was there.

There is no need to detail here how meticulously Kepler checked his rules against the observations of Tycho Brahe, whom he had served as an assistant before succeeding him as imperial mathematician at Prague. He was fully aware of the importance of empirical or observational

(Continued on page 25)



Left: The equal shaded areas indicate that a planet would move from A to B in the same amount of time as from C to D, or E to F.

Right: The table shows that for each planet the distance from the sun, cubed, is equal to the period squared, in agreement with Kepler's third law.

KEPLER'S HARMONIC LAW

Planet	Distance A. U.	Period Years	Distance Cubed	Period Squared
Mercury	0.387	0.241	0.058	0.058
Venus	0.723	0.615	0.378	0.378
Earth	1.000	1.000	1.000	1.000
Mars	1.524	1.881	3.54	3.54
Jupiter	5.20	11.86	141	141
Saturn	9.54	29.46	868	868
Uranus	19.18	84.01	7,060	7,060
Neptune	30.06	164.79	27,160	27,160
Pluto	39.52	248.43	61,720	61,720

# OBSERVING THE SATELLITES

## MAN IN SPACE

AMERICA'S space effort passed another milestone with the suborbital downrange ride of Comdr. Alan B. Shepard, Jr., in a Mercury capsule on May 5th. The flight of "Freedom 7" lasted only 15 minutes from its take-off at 14:34:13 Universal time from Launch Complex Five at Cape Canaveral, Florida, to its splashdown near latitude 27° north, longitude 76° west, in the Atlantic Ocean. During the flight, which attained a height of 117 miles and a range of 302 miles, the 2,250-pound package reached a top speed of 5,160 miles an hour, relative to a nonrotating earth.

Primarily, this test shot was intended to study the astronaut's reactions, and secondarily to check the fitness of the capsule. Shepard's physiological responses and ability to perform certain tasks had been evaluated under laboratory conditions prior to the flight. Thus his performance in the actual mission would show how reliable these tests were, and how effectively they prepared a man for the real experience. Early reports state that Shepard's flight has validated the testing and training procedures. The National Aeronautics and Space Administration claims that the Shepard flight was the first in which the pilot assumed control of the orientation of his spaceship.

The booster for this Mercury ascent was a Chrysler-built Redstone missile. An earlier version had been used as the first stage of the Jupiter-C that placed Explorer I in orbit for America's first successful satellite launching. The enlarged Redstone is about six feet in diameter, and with its manned capsule and escape tower has an over-all height of 83 feet. The total lift-off weight is 66,000 pounds, and the initial thrust amounts to 78,000 pounds.

Among the many safety features in Project Mercury equipment is a special system designed to end a mission by pulling the capsule away from sudden trouble. If the direction or amount of the rocket's thrust is wrong, or if, for example, electrical power is lost, this emergency system notes the fact automatically. When this happens, or if the ground station or pilot senses trouble, the booster's thrust is terminated, the spacecraft is separated, and the escape rocket mounted on trusswork atop the capsule fires to lift the capsule at least 2,000 feet clear of the booster.

The complex design of the Mercury capsule is the result of more than two years of intensive development by McDonnell Aircraft Corp. engineers. Several thousand subcontractors and suppliers have also contributed to the high-priority project. Much of the capsule's hardware had to be redesigned to meet both the severe weight limitations imposed by

available boosters, and the need to assure the astronaut's safety.

Outwardly, the capsule's appearance has changed little during development: a truncated cone (the cabin) is capped by a short parachute canister, and this in turn is topped by another truncated cone that serves as an antenna cover. Only nine feet tall, the capsule is 74 inches in diameter at its blunt end. Yet this small package contains an environment for the passenger's survival, with an alternative system should any component fail. The main systems are entirely automatic, while the emergency ones are either automatic, activated by ground command, or controlled by the astronaut.

The construction of the cabin walls illustrates some of the problems that faced the designers. The pressure-tight inner vessel is made from two very thin layers of titanium, the outer of these corrugated for strength. Welding this assembly required the development of a


new technique. To the outside of the pressure vessel was welded a framework of titanium stringers, and the whole was carefully insulated to break the flow of heat to the cabin's interior. Then an overcoat of corrugated metal was fastened to the stringers, in order to withstand the heat and stress of re-entry. In Shepard's flight, the exterior layer used beryllium shingles, and a massive beryllium heat shield covered the skin at the blunt end. In aerodynamic flight, this end swings forward, regardless of the initial orientation, and takes the brunt of the heat impact. In later missions, an ablative shield will be used, whose vaporization at re-entry will cool the remaining structure.

A primary automatic control system serves to regulate the spatial orientation of the capsule during its orbital flight, as well as just before the retro-rockets fire, and during the final plunge into the atmosphere. Deviations from the desired orientation are detected by infrared sensors and gyros, and are corrected by hydrogen peroxide jets. In another mode of operation, the pilot can assume con-



Comdr. Alan B. Shepard, Jr., on the aircraft carrier U.S.S. Champlain, stands in front of the Mercury capsule that carried him to an altitude of 117 miles. The capsule's landing cushion, between cabin and heat shield, is held partly open by blocks behind Shepard's feet. NASA photograph.





The flight sequence of a Redstone rocket and its Mercury capsule is indicated in this National Aeronautics and Space Administration diagram of an early flight plan. Heights and distances are given in statute and nautical miles; 100 nautical miles equal 115 statute. After the Redstone booster ceases firing, the superstructure atop the rocket blasts away. This is a safety system which can pull the capsule away from the Redstone in case of emergency. The capsule then separates; its orientation and the rocket's are shown for the remainder of the flight.

trol in either of two ways. In one, his control stick transmits electrical signals to six of the 18 peroxide jets; in the other, the same stick mechanically actuates throttle valves in the thrust system.

The control stick is in the pilot's right hand as he sits strapped in his contour couch against the blunt wall of the cabin. Pressure around him is maintained with oxygen at five pounds per square inch. Should cabin pressure be lost, his suit will automatically inflate to this pressure. Carbon dioxide removal and humidity control are automatic.

During Shepard's flight, duplicate communications channels were provided and tested. Periodically he radioed to flight monitors reports which included the voltage of the battery power supply, the amount of hydrogen peroxide fuel remaining, the oxygen pressure in the cabin, the accelerative stress, and other details.

The maximum acceleration during Shepard's launching amounted to six times gravity, and then for 4½ minutes he experienced weightlessness. While the retro-rockets were not needed to initiate re-entry during this suborbital flight, they were tested, and found to develop 0.05 times gravity. (From orbital motion at

17,500 miles per hour, only a two-percent reduction in speed would be needed to cause re-entry.) Then, during downward return through the atmosphere, the acceleration mounted toward 11g, but the total time it exceeded 5g was only about half a minute.

The only reported malfunction during Shepard's flight was failure of a green signal light to show that the retro-rocket package had been jettisoned after firing. He pressed an override button that could release this package, and then the light came on, to signify that the event had already occurred on schedule.

One test planned was the recognition of geographical features with the periscope display on the instrument panel. These were seen, together with cloud cover. Later, the astronaut closed his eyes and touched many preselected controls on the complex instrument panel before him, in order to test his psychomotor reactions. Throughout the flight, cameras monitored the entire instrument board, as well as Shepard's actions and appearance.

The success of the May 5th experiment means that manned orbital flight in a Mercury capsule now awaits only improved performance of the Atlas booster. Meanwhile, additional suborbital Red-

stone flights by astronauts are scheduled.

Previously, the Atlas had been used four times in the Mercury program. On September 9, 1959, a Big Joe Atlas successfully sent up a "boilerplate" capsule, in order to test re-entry performance and recovery techniques. Next, on July 29, 1960, the Mercury-Atlas 1 (MA-1) test failed, due to booster malfunction. Later, on February 21st of this year, MA-2 successfully tested the capsule, but on April 25th MA-3 was blown up by the range safety officer when the booster did not perform as intended.

If we also count earlier satellite launchings, Atlas has already been applied 13 times, 12 of these with superstructures attached. In the latter firings, only four missions have been completely successful, though very few of the failures were catastrophic malfunctions. Attachment of the Mercury to the Atlas will be further tested before a manned launching.

The main objective of the Mercury program is to carry a man three times around Earth and land him safely. Modified capsules are being developed for longer-duration animal flights, mainly to study improvements in the life-support systems. At the same time, the U. S. Air Force is developing Dyna-Soar, a 5-ton

delta-wing glider that will attempt to carry a man into orbit atop a Titan II rocket. The winged design is expected to give better control over re-entry heating than the ballistic-parachute system of Mercury does.

Advances in manned space exploration beyond Mercury and Dyna-Soar are sure to be very costly. Recently, Congress has been asked to increase NASA's funding for the fiscal year 1962 to \$1,784,300,000. Much of the increase is earmarked for Apollo, which is a successor program to the man-in-space effort, and for expanded research in biological problems related to space travel. Project Saturn, which will supply the booster for Apollo flights, already was the biggest item in NASA's budget.

Large as these funds for 1962 are, they are small compared to the projected cost in years ahead. Annual expenditures in excess of four billion dollars will be asked, and some 40 billion may be required before the first U.S. expedition sets foot on the moon. So great is the effort required for this development that it may be said that the only component now nearly ready to go is man himself.

#### VOSTOK FLIGHT RECORDS

THE Fédération Aéronautique Internationale, with headquarters in Paris, France, has established rules under which world records of various kinds may be claimed for manned spaceflights. Immediately after Yuri Gagarin's 108-minute journey in Vostok on April 12th, claims were filed for duration, altitude (203 miles), and total load (4,725 kilograms) — all in the class of manned orbital flights.

Nearly seven weeks after the event, certain facts required by the rules for substantiation of the claims were submitted to the federation, and thus made public for the first time. According to unofficial translations of Tass reports, Vostok was launched by a rocket with six engines. The trip began at Baikonur, near the Aral Sea, at latitude 47° north, longitude 65° east, and is said to have ended near the village of Smelovka, Ternov district, Saratov region.

These locations confirm the deductions cited on page 330 of last month's issue for the beginning and end points of the flight.

MARSHALL MELIN

Research Station for Satellite Observation  
P. O. Box 4, Cambridge 38, Mass.

#### CORRECTIONS

On page 330 of the June issue, first paragraph, the longitude at which Vostok passed northward over the equator should read 15° east, not 15° west. This error was noted by Paul K. Hoover, Salt Lake City, Utah.

On page 331, the right-hand picture of Explorer XI's gamma-ray telescope is inverted.

## Astronomers Discuss Project West Ford

TO provide reliable intercontinental radio communications, W. E. Morrow, Jr., of Lincoln Laboratory, has proposed placing into orbit around the earth a belt of numerous tiny metallic needles. Already plans have been made to launch a test belt of such dipoles, 75 pounds of them to be dispensed from an artificial satellite to form a cloud that will have spread completely around the earth after a month. This program, now known as Project West Ford, was briefly described in *SKY AND TELESCOPE*, December, 1960, page 329.

The question of whether West Ford belts will interfere with astronomical observations has aroused much discussion, and four articles in the April *Astronomical Journal* deal with this problem. Dr. Morrow and D. C. MacLellan report on some of the properties of the proposed belts. The first test belt, they predict, should cause quite limited sky brightening and its effect on radio astronomy measurements would be very small. However, later West Ford belts may contain much larger quantities of the tiny needles.

William Liller of Harvard Observatory points out, "Optical astronomers must carefully evaluate the effects of the orbiting dipoles in the test belt as accurately and as quickly as possible so that an upper limit can be set on the size of future experiments, if any." He recommends photographic observations with fast, wide-field cameras to record the belt, which during its first two months aloft would appear as a very faint, narrow, luminous streak spanning the sky. In addition to photoelectric and polarimetric studies, he stresses the value of visual observations, for example naked-eye sightings through a slowly rotated sheet of polaroid.

Calculations by A. E. Lilley of Harvard confirm the Morrow-MacLellan appraisal that the test belt will have negligible

effects on broad-band radio astronomy observations, but there may be disturbances to radio spectrum-line studies. Later operational belts, densely sown with dipoles, will create unavoidable problems for radio astronomers. Dr. Lilley adds the following broader comment:

"The threat to the conduct of radio astronomical research by orbiting media should be viewed in perspective with other sources of man-made interference. New communications and navigation systems employing powerful transmitters in earth satellites will be soon forthcoming. These technological advances pose another source of severe interference to radio research. The pursuit of basic science and the progress of space radio technology represent needs of man which must be advanced. For the impending interference a simple solution exists: allocation of clear frequency bands for basic science. This action is imperative and must ultimately rest on national and international agreements."

#### OBSERVING TIME ON KITT PEAK

The national observatory on Kitt Peak in southern Arizona was established to provide American astronomers with modern observing instruments. The director, N. U. Mayall, has announced a policy allocating about 60 per cent of the observing time to visiting astronomers and qualified graduate students.

To supplement the 36- and 16-inch reflectors currently operating on the mountain, observatory headquarters in Tucson contain laboratory equipment that includes a digital computer, a microphotometer, measuring engines, and a spectrocomparator. Additional information is available from Kitt Peak National Observatory, 950 N. Cherry Ave., Tucson, Ariz.

#### JOHANN KEPLER AND THE LAWS OF PLANETARY MOTION

(Continued from page 22)

proof in astronomy. But there was another side to Kepler, more difficult for us in the 20th century to understand.

He was a metaphysician and mystic, whose mind was filled with the lore of Plato, Pythagoras, and Plotinus. Thus, while he laid the foundations of celestial mechanics, all his life he was entranced by the hope of finding a relationship between distances of the planets from each other and the ancient doctrine of the five regular solids. He tried to establish a connection between planetary orbits and the rules of musical harmony. Indeed, his passion for the harmony of the spheres probably first attracted him to astronomy and kept him there despite the practical difficulties of living in the midst of tremendous political and religious chaos.

A further aspect of Kepler's complex personality deserves mention. He was not eaten up by a passion to be the first to make a scientific discovery, and did not have the secretiveness characteristic of some brother astronomers. Instead, he sought co-operation among astronomers, for example in observing the total solar eclipse of October 12, 1605. In an age of religious controversy, he was without bigotry. Although a Lutheran, he sided with the Calvinists on some theological points, and at the same time remained on good terms with the Roman Catholics. When Jesuit missionaries at the court of the Son of Heaven in Peking failed to persuade Galileo to send them eclipse calculations, Kepler immediately supplied a part of the Rudolphine Tables which he was compiling. The fact was that Kepler's vision of the harmony of the universe raised him above the strife of his time.

# Amateur Astronomers

AN ACTIVE AMATEUR IN ST. PAUL

**D**URING the past 25 years, I have devoted an increasing amount of time to studying and teaching astronomy. My first telescope, completed in 1937 when I was 14, initiated a succession of home-constructed instruments, and recently I built my Cobb Road Observatory.

Although today's trend is toward compactness, this observatory is a bit larger than originally planned. A dome 16 feet in diameter sits at one end of the 21-foot structure, leaving a rectangular area five by 16 feet for instrument storage, tables, and chairs.


The truncated-cone shape of the dome was chosen for simplicity of construction;  $\frac{1}{2}$ -inch tempered Masonite was wrapped around the 1-by-4-inch frame ribs. It was covered with tar paper, with aluminum roofing paint for better heat reflection.

The dome does not turn on a track; it swivels on 32 roller-skate wheels mounted around the dome well on the observatory roof. Ten more wheels set horizontally on a flange around the outside of the well prevent the dome from slipping. So far I have turned it by hand, as the rolling friction is quite small, but I plan to install a crank or a motor. The observatory's interior is being finished in light blue, with the inside of the dome a flat black. There is ample wiring for red and white working lights.

At present the building is filled with all kinds of telescopic equipment. A  $4\frac{1}{2}$ -inch  $f/12$  reflector has a 7-inch tube to give maximum side clearance for reducing thermal effects on the incoming light beam. This instrument splits the "double double" in Lyra very easily at a magnification of 250. An array of medium-sized reflectors includes a solar

telescope with an unsilvered antireflection-coated 6-inch mirror, and 6-inch  $f/12$  and 8-inch  $f/8$  telescopes.

In the center of the observing area is my 125-pound, 12-inch off-axis reflector. Its 196-inch focal length gives a primary image of the moon  $1\frac{1}{4}$  inches in diameter; low-power views promise to be excellent,



The dome of the Cobb Road Observatory in St. Paul, Minnesota, has an opening five feet wide that allows the building to reach outside air temperature quickly, as well as providing an ample view of the sky for observing.

as soon as astigmatic corrections are finished. Astronomical photography has always been one of my major interests, and in my equipment are three star cameras, between seven and 40 inches focus. The 12-inch telescope will be used for visual and photographic observation of the moon and planets.

About seven years ago I began teaching a night course in astronomy at Macalester College, and four years later took over the day course. This class's observation of the November 7, 1960, transit of Mercury was described in *SKY AND TELESCOPE* for January, 1961. Now in the process of expanding, the college is building a new science hall, with planetarium and observatory.

My profession, optometry, seems to be fading into the background as more time

is spent on astronomical activities. Last January I began a series of 22 weekly half-hour local television presentations, entitled "Astronomy Today: A Guide to Other Worlds in Space." The programs emphasized the solar system, giving planetary facts and theories and observing hints.

The local Moonwatch team, which I am currently leading, meets at the Cobb Road Observatory, and star parties take up the weekends. We prefer groups of about 25 persons, for each guest can then move freely about the building. However, one night 60 adults and cub scouts attended an open house, which was highlighted by a passage of Echo I over our area.

SHERMAN W. SCHULTZ, JR.  
944 Cobb Rd.  
St. Paul 12, Minn.

## MIDDLE EAST CONVENTION

Members of the Astronomical League's Middle East region will convene at Hershey, Pennsylvania, on Saturday, August 19th. The meeting is being organized by the Astronomical Society of Harrisburg.

Lectures, exhibits, papers, and round-table discussions are scheduled to fill the day-long program, highlighted by the traditional banquet. Inquiries should be sent to Edward L. Naylor, 320 Wilhelm Rd., Harrisburg, Pa.



Just after evening fell on November 21, 1960, Sherman Schultz, Jr., took this photograph of his Cobb Road Observatory with a spectacular celestial backdrop. Over the doorway are the bright planets Venus (left) and Jupiter. Above them is the thin crescent moon, its night side faintly illuminated by sunlight reflected from the earth. This home-built St. Paul, Minnesota, observatory is a local center for the popularization of astronomy.



## REPORT FROM ROCHESTER

IT WAS lilac festival weekend in Rochester, New York, when the Northeast region of the Astronomical League held its annual get-together. Although skies were cloudy on Friday evening, May 26th, more than 50 persons enjoyed the hospitality of the George Keenes and inspected their back-yard observatory (featured in the January, 1959, SKY AND TELESCOPE).

Saturday morning the delegates visited Bausch and Lomb Optical Co. to watch the pouring of optical glass. The first session at the Manger Hotel included papers on topics ranging from antique telescopes to activities of the Rochester Moonwatch team.

After lunch, Edgar Everhart, Mansfield Center, Connecticut, showed color slides of the September 17, 1960, solar halo display (January, 1961, issue, page 14). A feature of the eight-paper session was the exhibition of a clock giving standard and sidereal time, positions of the sun and moon, and lunar phases. This instrument was built by Edward Root of Rochester as a cloudy-weather project.

At the business session, Mrs. Margaret Tumiki, New London, Connecticut, was elected Northeast chairman, and Parker Atwood, East Haven, Connecticut, vice-chairman; Walter Wyman, Batavia, New York, was re-elected secretary, and Jack Welch, Springfield, Massachusetts, treasurer. Springfield was announced as the site for the 1962 convention, and the 1963 meeting is scheduled for Stratford, Connecticut.

Highlighting the banquet were solar-prominence movies filmed and narrated by Walter Semerau of Kenmore, New York.

## AN AMATEUR OBSERVATORY IN THE TRANSSVAAL

A successful star party marked the public opening of C. N. Williams' new observatory, located at Bedford View near Johannesburg, South Africa. Many amateur astronomers and friends from Johannesburg, Germiston, and Boksburg came to admire the observatory building and the well-constructed instrument it houses.

Dr. Williams' 5-inch refractor and its two finders are mounted equatorially on a short iron column bolted to a concrete block 16 inches in diameter. The telescopes are driven by a small synchronous motor.

The rectangular brick observatory has a sliding roof of corrugated steel, which moves on rails supported by piers on the building's south side. It is large enough to hold 20 people easily, and is fitted with a table, chairs, a library, and a radio receiver for time signals.

Dr. Williams was the team leader of the Johannesburg Moonwatch, which contributed many observations of early artificial satellites. He now plans to use his

telescope for regular observation of the moon's surface features, a study for which his refractor is well suited and which has been neglected by many amateurs here in South Africa.

J. H. BOTHAM  
94 Ascot Rd.  
Judith's Paarl

Johannesburg, South Africa

## LAKELAND, FLORIDA

Twenty-five amateurs, many formerly associated with other astronomical organizations, are members of the new Central Florida Astronomical Society. Adults and juniors interested in joining the group should contact Carson H. Sammons, 1810 Fredericksburg Ave., Lakeland, Fla., MU 2-3223.

## SOCIETY LISTING

The October issue of SKY AND TELESCOPE is tentatively scheduled to carry Here and There with Amateurs, a listing of astronomical societies whose membership is open to the public. Any group that wishes to be included should send to this magazine by August 15th a post card stating: the organization's name, the city in which it is located, and the name, address, and telephone number of an official who can furnish details of the group's activities.

In addition, please answer the following questions by number:

1. Is the organization a member of the Astronomical League?
2. Does the society belong to the Western Amateur Astronomers?
3. Is there a junior section?
4. Is this an independent junior group?
5. Does the society include a subscription to SKY AND TELESCOPE as a privilege of membership?

To insure an up-to-date listing, each club should submit the post card promptly; otherwise the information published in the 1960 Here and There with Amateurs (October issue, page 209) will be repeated.

## WESTCHESTER ASTRONOMERS LEAGUE

Thirty-five amateurs from all areas of Westchester County, New York, attended the first meeting of the Westchester Astronomers League. The gathering was addressed by Edgar Paulton, past president of the Amateur Astronomers Association of New York, and by Frank L. Aime, who gave an illustrated lecture on the home-made observatory and 16-inch Cassegrainian telescope that he built. The new group will meet regularly at the Hudson River Museum in Trevor Park, North Yonkers, on the third Friday of each month at 8 p.m. All are welcome.

VICTOR F. VOLK  
749 North Broadway  
Hastings-on-Hudson, N.Y.

## THIS MONTH'S PROGRAMS AND CONVENTIONS

Detroit, Mich.: Astronomical League, Henrose Hotel. July 1-3.

New Orleans, La.: Pontchartrain Astronomy Society, 8 p.m., Metairie playground auditorium. July 7, M. O. Guidry, Texaco, Inc., "Use of Setting Circles," and Richard Sarradet, De La Salle High School, "Generalities of the Electromagnetic Spectrum."

Shreveport, La.: Shreveport Junior Astronomical Society, 7:30 p.m., Centenary College science hall. July 15, Ted Gangl, Texas Astronomical Society, "The Sky Is Not the Limit."

## STELLAFANE MEETING

The 27th Stellafane convention will be held August 12th, on Breezy Hill at Springfield, Vermont. This first Saturday after new moon in August was chosen by popular vote of the 100 hardy amateurs who gathered under the big tent during the torrential downpour that washed out last year's meeting.

Edgar Everhart of Mansfield Center, Connecticut, is to be master of ceremonies for the program, which includes a workshop session on mirror polishing and figuring, a meeting of the Maksutov Club, the regular telescope contest, and observing. Twilight talks will include "Deep-Sky Wonders," by Walter Scott Houston, Henry Specht's "Photographic Measurements of Variable Stars," and "Solar Research" by Walter Semerau.

JAMES W. GAGAN  
17 Bellevue Ave.  
Revere 51, Mass.


## VARIABLE STAR SOCIETY IN GERMANY

In 1950 the Berliner Arbeitsgemeinschaft für Veränderliche Sterne (BAV) was founded by a group of amateur astronomers in Berlin especially interested in variable stars. The first such organization in Germany, it soon became a center for systematic observing by amateurs. We also co-operate closely with the AAVSO, sending them monthly reports.

Our headquarters are located in the public Wilhelm Foerster Observatory, where amateurs carry on all the scientific work, the institution supplying many of the necessary instruments. The most effective one for our purposes is the 12-inch Bamberg refractor.

Some 20 members of the BAV are now actively engaged in timing minima of eclipsing variables. Several times a year they receive the BAV Rundbrief, a circular containing about 12 pages of observations, ephemerides, and other pertinent information. We now have 75 eclipsing stars under systematic observation; these were listed in the Rundbrief for January, 1958.

When most of our members lived in Berlin, it was unnecessary to construct



Reproduced almost full size, this combination chart for U Cephei was prepared by the Berlin Group for Variable Star Observing. Seventy-five such diagrams, covering all stars in the society's program, will soon be available.

field charts for these stars, since the star charts and atlases at the observatory could be used. However, now that our membership has expanded, we are mapping the fields of all stars in our regular program.

A typical sheet is approximately six inches wide by 8½ inches high, and contains a constellation sketch and three progressively more detailed maps of the variable's region. Also included is information about the star's co-ordinates, magnitudes at maximum and minimum, duration of eclipse, and period. Other variables in the same field are marked,

to prevent their accidental use as comparison objects. Observers interested in eclipsing binaries may buy these charts for 10 cents each.

Our organization would like to find collaborators in the United States. There is a special need for observers with large telescopes (12-inch or so) for work on the fainter eclipsing variables.

**WOLFGANG QUESTER**  
BAV, Wilhelm Foerster Sternwarte  
Papenstrasse 2  
Berlin-Schöneberg  
West Germany

### \*\*\* AMATEUR BRIEFS \*\*\*

Much-enjoyed activities of the Clinton County Astronomy Club in Frankfort, Indiana, have been field trips to various places of astronomical interest. At 5 o'clock one early spring morning, club members climbed into the back of a panel truck, furnished with hay bales, for a trip to Chicago and the Adler Planetarium. The expedition was successful, although all the way home the truck unwittingly played tag with a tornado, at one time being only five minutes ahead of the storm's center!

Reports of observations received by the Montreal Centre of the Royal Astronomical Society of Canada are filed in a pigeonhole rack over the library desk and then picked up promptly by the respective section chairmen.

Suggesting summer uses for a telescope, the bulletin of the Amateur Astronomers Association of New York City quotes from a letter by the old-time telescope maker, John Brashear: "Of an afternoon, Dr. Samuel Langley would come over to my shop to watch the baseball games as the park was in sight and a telescopic view of the game was very satisfactory."

Dewey Schwartzberg, Jr., 14, vice-president of the Shreveport Junior Astronomical Society, won top honors at the Louisiana State Junior Science Fair, and will represent his state at the national fair to be held in Kansas City. In his project, the Milky Way and the Messier objects he had personally observed were represented on a globe.

A 100-volume astronomical library gathered by the amateur Alvin J. Aucoin of Houston, Texas, before his death in 1940, has been donated to the local Lamar High School Astronomy Club by his wife and son. In the collection are several 19th-century works, by such writers as Grant, Smyth, Proctor, and Loomis.

One member of Rochester Moonwatch, according to the team's leader, Russell Jenkins, prepared diligently to observe satellite 1960:1. He computed the precise time and altitude of an expected meridian transit, arranged his equipment, and then spent several minutes adapting his eyes to the darkness. At the predicted time, 1960:1 appeared, a brilliant, unmistakable object. Much to the Moonwatcher's chagrin, he realized his elaborate preparations were for Echo II!

New quarters of the Astronomical Society of South Australia are located at the Norwood Boys' Technical High School, Adelaide. Recently, a used 12-foot observatory dome was donated to the society by Dave Rumball. Transporting the dome to the school grounds proved quite an undertaking, but with 13 good men, a large flat-bed truck, and an escorting convoy of a dozen cars, it was successfully managed. The dome is now being repaired, scrubbed, and painted, and its concrete foundation built. G. B. C.

# Ten Years of Solar Eclipses

**D**URING the decade from January, 1961, through December, 1970, the shadow of the moon falls upon the earth 22 times. These events occur as eight total, eight annular, and six partial eclipses of the sun. Every continent will see at least one of them, weather permitting.

The tables contain some information on the dates and locations of eclipses in this period. On the following pages is a map indicating the central lines of the total and annular events.

Although the number of eclipses is the same as in preceding and succeeding decades, eight of them are total during



Four saros cycles before February 15, 1961, the eclipse of January 1, 1889, was observed along a path running from north of San Francisco to the Hudson Bay region. This photograph was taken by the Harvard expedition in Willows, California, with a 13-inch telescope.

## PARTIAL SOLAR ECLIPSES 1961-1970

Date	Magnitude	Where Visible
1964 Jan. 14	0.559	Antarctica, Tasmania, S. America
Jun. 10	0.754	Australia, Tasmania, New Zealand
Jul. 9	0.322	Asia, Canada, Arctic
Dec. 3-4	0.752	Hawaii, Japan, Asia, Alaska
1967 May 9	0.720	N. America, Greenland, Scandinavia
1968 Mar. 28-29	0.899	Antarctica, S. America

The magnitude of an eclipse of the sun is the fraction of the solar diameter covered by the moon at maximum phase, measured along the line connecting the centers of the disks.

1961-70, compared with six in 1951-60, and seven in 1971-80.

Three total eclipses are of special interest because of the areas over which they pass. The first occurred last February, crossing southern Europe and part of Asia (SKY AND TELESCOPE, April, 1961). It will probably prove to be the most widely observed eclipse of the period.

On July 20, 1963, the moon's shadow

will race across North America. Although much of the path will be over land, the short length of totality may hamper some types of observing programs. In Alaska, the duration will be about 1½ minutes, but totality will have dwindled to only 59 seconds by the time the shadow reaches the coast of Maine.

The somewhat longer totality associated with the eclipse of March 7, 1970, makes

it more favorable. The path crosses Central America and Florida and skirts the eastern seaboard. Although totality is largely over water, the accessibility of land observing sites, combined with the high altitude of the sun (totality occurring shortly after noon in the United States), is likely to attract many professional and amateur astronomers. The maximum duration of totality is 3½ minutes, near Coatzacoalcos in southern Mexico, but it will last about three minutes in Georgia and the Carolinas.

The November 2, 1967, eclipse is unique among the 22, because the axis of the moon's shadow will not touch the earth's surface. Sweeping past the Southern Hemisphere, the edge of the umbral shadow will fall on a small area of ocean north of the Weddell Sea, but the center of the shadow cone will miss the earth by about 14 miles. Such a near miss last took place on October 23, 1957, in the same part of the Antarctic. This total eclipse will probably be observed by fewer people than any other during the decade.

Occurring in high northern latitudes, the eclipse of September 22, 1968, will follow a north-to-south path, rather than the more common west-to-east track. The eastward sweep of the shadow will be nearly matched by the turning of the earth, resulting in a central line that ends slightly west of where it started.

The longest duration of totality will come with the eclipse of May 30, 1965. Unfortunately for observers, the central line falls almost entirely over water. The shadow passes north of New Zealand, crosses the Pacific, and leaves the earth in Peru, not from from the coast. In mid-Pacific the sun will be hidden for a total of 5½ minutes.


South Americans who miss the 1965 eclipse because of the shortness of its path over the continent will be solaced by another passage of the moon's shadow

## TOTAL AND ANNULAR SOLAR ECLIPSES 1961-1970

Date	Type	Duration m s	Width Miles	Location
1961 Feb. 15	T	2 45	164	Europe, Asia
Aug. 11	A	6 35	323	South Atlantic
1962 Feb. 4-5	T	4 08	91	Central Pacific
Jul. 31	A	3 33	100	Atlantic, Africa
1963 Jan. 25	A	1 01	46	South Atlantic
Jul. 20	T	1 40	63	North America
1965 May 30	T	5 16	124	Pacific
Nov. 23	A	4 05	108	Southern Asia
1966 May 20	A	56	40	Africa, Asia
Nov. 12	T	1 57	52	S. America, S. Atlantic
1967 Nov. 2	T			Weddell Sea
1968 Sep. 22	T	40	68	North Central Asia
1969 Mar. 18	A	1 14	44	Indian Ocean, Pacific
Sep. 11	A	3 15	104	North Pacific
1970 Mar. 7	T	3 28	98	North America
Aug. 31- Sep. 1	A	6 48	188	South Pacific

Successive columns give the date of each eclipse, its type (total or annular), the maximum length of time between second and third contacts, the maximum width of the path from which the eclipse may be seen as total or annular, and the general area of the earth over which the path passes.





Paths of total eclipses (solid lines) and annular (dashed lines) occurring in the decade 1961 to 1970, drawn on a U. S. Hydrographic Office chart. R, N, and S indicate local sunrise, noon, and sunset, respectively.

only a year and a half later, on November 12, 1966. This time the eclipse will go slightly south of the end of the previous one and then continue across Bolivia, Argentina, and southernmost Brazil.

The only total eclipse with a central line that does not touch any continent will come on February 4-5, 1962. Because

the path crosses the international date line the event will occur on both the 4th and 5th.

Two of the annular eclipses are quite lengthy. On August 11, 1961, six minutes and 35 seconds will elapse between second and third contacts. Even this is 13 seconds less than the duration predicted for the

last eclipse of the decade, occurring on August 31 and September 1, 1970. But the central line of the latter falls over the Pacific and the former crosses the South Atlantic, where few people will view the spectacle of the bright solar ring around the moon.

There will be ample opportunities for



The hatched area in the Weddell Sea indicates shadow coverage for the noncentral total eclipse on November 2, 1967. A small part of the path in the Arctic of the September 22, 1968, eclipse is omitted.

such observations, however, since annular eclipses will be visible from Africa, Europe, Asia, and South America. Although not as important to astronomers as total eclipses, because the corona is not seen, annular eclipses draw much local attention and are useful in geodetic studies.

Two successive annular eclipses, July

31, 1962, and January 25, 1963, may be valuable for such purposes. Since each crosses parts of two continents, measurements may be made to determine transoceanic distances, thus tying together trigonometric surveys of the two land masses.

In all, the decade should be of interest

to eclipse observers, since the paths of three total eclipses are largely over land, with three others presenting reasonably good observing possibilities. *SKY AND TELESCOPE* will carry more detailed information about some of the coming eclipses near the times of occurrence.

R. N. W.

## Deep-Sky Observers



An extremely fine, imported Zuiho binocular with a wide field of view—10½°!

**\$35.00**  
Tax Paid

Complete with beautiful case. Delivered anywhere in the Continental United States.

**AMERICAN TYPE WIDE ANGLE**  
7 x 35 Central Focus

### SPECIFICATIONS

Magnification, 7x. Objective aperture, 35 mm. Eyepiece aperture, 24 mm. Exit pupil, 5 mm. Field of view at 1,000 yds., 551 ft. Field-of-view angle, 10.5°. Height, 128 mm. Breadth, 195 mm. Net weight, 950 grams. Weight of pigskin case, 400 grams.

We have sold hundreds of pairs of these binoculars with the utmost satisfaction. They are unconditionally guaranteed. If not pleased, return them and your money will be immediately refunded.

Established 1936



## ADIRONDACK RADIO SUPPLY

185-191 West Main St.

P.O. Box 88, Amsterdam, N. Y.

Ward J. Hinkle, W2FEU — Owner

## BRANDON OCULARS



These orthoscopic oculars are the choice of the amateur as well as the professional astronomer. Seven sizes available: 4-mm., 6-mm., 8-mm., 12-mm., 16-mm., 24-mm., and 32-mm. focal lengths. Standard 1¼" outside diameter.

**\$15.95 each**

Brandon Oculars may be purchased at —

Cave Optical Co., 4137 E. Anaheim St., Long Beach 4, Calif.

Cleveland Astronomics, 7618 Lawn Ave., Cleveland 2, Ohio

American Science Center, 5700 North-west Highway, Chicago 46, Ill.

Optron Laboratory, Box 25, D.V. Station, Dayton 6, Ohio

Adler Planetarium, 900 E. Acheson Bond Dr., Chicago 5, Ill.

Polaris Telescopic Shop, 14319 Michigan Ave., Dearborn, Mich.

Star-Liner Co., 887 Sherbourne Dr., Inkster, Mich.

—or order direct. We will ship airmail.

## LIBRASCOPE P. R.

Guaynabo

Puerto Rico

Formerly BRANDON INSTRUMENTS

# OBSERVER'S PAGE

Universal time (UT) is used unless otherwise noted.

## MUTUAL PHENOMENA OF JUPITER'S SATELLITES

**D**URING the second half of this year, the orbits of Jupiter's four bright satellites will be turned nearly edgewise to the earth. Consequently, there will be a number of occultations and eclipses of one Jovian moon by another.

The accompanying list of predictions includes only the more conspicuous of these phenomena visible from the United States. It has been extracted from a full list in the 1961 *Handbook of the British Astronomical Association*.

The mutual occultations are best observed with large telescopes and high powers capable of revealing clearly the satellite disks. Careful timings of the contacts are desirable.

Smaller telescopes are well suited for following the eclipse of one satellite by the shadow cone of another. In this case, the decrease in brightness and subsequent recovery of the eclipsed moon can be observed by means of comparison with other satellites or with field stars. A particularly striking event of this class is the eclipse of II by III on September 27th, when the former will fade to near-invisibility and brighten again within a 54-minute interval. An observer wishing to derive a light curve of these rapid changes might well use a tape recorder for his brightness estimations.

1961	Occultation of	Universal Time
Aug. 29	II by III	7:04 - 7:11
Sep. 26	II by I	6:03 - 6:31
Oct. 4	II by III	1:41 - 2:09
Oct. 14	II by I	1:38 - 1:47
Nov. 7	II by I	22:36 - 22:39
Nov. 22	II by I	3:33 - 3:36

Note that the predicted durations are mostly only a few minutes, but in some instances much longer. Brief events result when one moon is on the nearer side of Jupiter, the other on the farther side, so they pass quickly in opposite directions. The very extended duration of the September 27th eclipse comes from II's slow motion just after eastern elongation, which nearly matches the speed of III and its shadow.

1961	Eclipse of	Universal Time	Mag.
Aug. 22	II by III	6:17 - 6:24	0.30
Sep. 12	II by I	2:49 - 3:21	0.57
Sep. 19	II by I	6:40 - 6:54	0.64
Sep. 27	II by III	4:36 - 5:30	0.98
Oct. 7	II by I	1:42 - 1:50	0.66
Nov. 8	II by I	0:45 - 0:49	0.38
Nov. 27	I by II	1:33 - 1:36	0.33
Dec. 4	I by II	3:47 - 3:50	0.33

Mag. is the magnitude of eclipse — the fraction of the diameter of the remoter satellite that is shadow-covered at mid-eclipse. The eclipses of November 27th and December 4th are annular, the smaller satellite being in front.

## SZ HERCULIS PREDICTIONS

The Algol-type variable star SZ Herculis is now conveniently placed for observation with telescopes of 6-inch aperture or larger. A finder chart and other information were given on page 508 of the June, 1960, issue. The magnitude range is 10.2 to 11.9, eclipses occurring every 0.818 days. Only 2.2 hours are taken for this variable to fade from maximum to minimum light, and an equal interval to recover full brightness.

Because the period of SZ Herculis is changing, new timings of its minima are very desirable. All that is needed is a series of magnitude estimates at roughly 10-minute intervals during the hour preceding and the hour following minimum. Readers who make such observations are invited to send copies of their records to SKY AND TELESCOPE for analysis.

During the next two months, minima occurring at convenient hours for North American observers are predicted for the following Universal times, expressed in hours and tenths:

July 1, 6<sup>h</sup>.1; 6, 3.9; 10, 6.1; 15, 3.9; 19, 6.1; 24, 3.9; 28, 6.0.

August 2, 3<sup>h</sup>.9; 6, 6.0; 11, 3.8; 15, 6.0; 20, 3.8; 24, 6.0; 29, 3.8.

Other minima may be predicted by adding multiples of 19.63 hours to these times.

## Astronomy Teaching Aids

### ESSCO Publications

Teachers of astronomy, both amateur and professional, will find the following ESSCO study materials useful for their classes. The test charts are challenging fun for the amateur astronomer, too!



- SC1 Equatorial constellation chart — with star designations
- SC1T Test equatorial chart — without star or constellation names
- SC2 Circumpolar constellation chart — with star designations
- SC2T Test circumpolar chart — without star or constellation names
- S508A Ecliptic-based star map — with equatorial grid and names
- S508B Ecliptic-based star map — with equatorial grid, without names
- S508B Ecliptic star map list — positions and magnitudes for 224 stars
- S505 Nine-inch protractor on paper — for planet orbit drawings
- S511 Inner planet chart — orbits of Mercury, Venus, Earth, Mars
- S512 Outer planet chart — orbits of Mercury to Saturn
- S501A Special rectangular co-ordinate paper — for star maps
- S502 Polar co-ordinate paper — for circumpolar star maps
- S600 Aitoff's equal-area projection of the sphere — 13 inches wide

Price for each item listed above: 1 to 9 sheets, 10 cents each; 10 to 24 sheets, 8 cents each; 25 to 99 sheets, 6 cents each; 100 or more, 5 cents each.

**SPECIAL: DC5** — Large wall chart of the Draper spectral classes, printed in four colors and mounted on canvas. Limited quantity. (Slightly damaged along edges.) **\$14.00 each**

**SKY PUBLISHING CORPORATION**  
Cambridge 38, Massachusetts





Left: Mr. Herring's drawing of Lassell was made with a 12½-inch reflector at 275x, on August 1, 1960, at 3:30 UT. South is above, east to the right. The dome mentioned in the text is about one inch from the picture's right edge. Right: Part of sheet D5-a of G. P. Kuiper's "Photographic Lunar Atlas," from a Mount Wilson 100-inch reflector plate.

#### OBSERVING THE MOON — LASSELL

**N**EAR the western border of Mare Nubium is the small crater Lassell, east of Alpetragius and nearly midway between Davy on the north and the Straight Wall on the south. Approximately 14 miles in diameter, it lies near the center of an almost obliterated ring, one of a chain of large ancient rings that paralleled the western edge of the mare (SKY AND TELESCOPE, April, 1959, page 335).

The crater is an irregular hexagon with walls that are generally low and dilapidated. The highest portion of the rim may be about 2,300 feet, on the west, and a peak of somewhat lesser height sits on the east. The outer wall drops sharply to the level of the outside plain on the east, but the western slopes are much more gentle. Narrow gaps interrupt the walls on the north and south, with a number of small craters near these openings.

Inside Lassell there are few conspicuous details except for a group of low hills and ridges scattered about the floor. I have suspected an extremely delicate rille concentric with the west wall, a feature that other observers should look for. The inner east wall has an easily seen terrace, similar to those in numerous craters on the moon. Probably they were formed by partial collapse of the walls when support on the inside was withdrawn; it may be that subsidence has played an important part in crater formation.

The plain surrounding Lassell contains a number of low ridges, mostly oriented north and south. They appear similar to other wrinkle ridges, many of which tend to be concentric with mare borders. These small ridges require oblique lighting to be well seen. In the Lassell area are also numbers of craterlets, ranging from small pits scarcely a mile across to 6-mile

Alpetragius B, prominent in the left part of the accompanying drawing.

East of Lassell is a clump of low irregular hills, one of these small peaks having a readily seen summit craterlet. Near these hills is a curious group of several coalesced craters, bounded on the east by a little hill and on the west by several saucerlike depressions. Adjoining these

craters on the south is a small circular hill. With a diameter of 3.7 miles and sides inclined less than three degrees to the horizontal, this gentle swelling has an over-all height of only about 500 feet. It seems very similar to domes occurring elsewhere on the moon's surface.

ALIKA K. HERRING

1312 Arlington Ave.

Anaheim, Calif.

## STAR-LINER COMPANY

Manufacturer of Reflector Telescopes to 20" — 15 Years Experience

Announcing NEW features in the STAR-LINER EQUATORIAL MOUNT:

1. Designed for stability and beauty. Cast aluminum, carries 6" to 12.5" telescopes.
2. Polar adjustments are made precisely with a fine-aligning device.
3. Entire mounting swings through 360° for quick alignment on the local meridian. Satellite tracking is made simple with this exclusive feature.
4. Tube rotates fully within cradle. 6" to 12.5" cradles are interchangeable, permitting use of same mounting with different tube units.
5. Steel shafts, 1.750" diameter, ride on brass oil-lite bushings (never need oiling).

Compare these features with other mountings on the market today. Two years in development, this Equatorial Mounting is unexcelled in actual performance.

### Newly Designed Mirror Cells — The ADJUST-A-CELL

This cell has the following advantages over all others: It is completely adjustable to the exact diameter of the tube; mirror and cell together can be moved 1¼" forward or backward in tube; and there are fine-collimation aligning screws and locking bolts for mirror tilt. (Patent applied for.)

Size	Price	Special Orders for:
4¼"	.....\$ 6.50	16" with 9-point flotation .....
6"	.....\$ 8.50	.....\$ 78.00
8"	.....\$12.50	20" with 12-point flotation .....
10"	.....\$19.75	.....\$142.00
12½"	.....\$27.50	


We carry a full line of quality parts and accessories. See these and complete STAR-LINER telescopes at the Astronomical League convention in Detroit, Mich., July 1-3.

A wealth of helpful information is yours on request. Write for our specially compiled catalog.

### STAR-LINER Company

887 Sherbourne Drive

Inkster, Michigan



### COMPLETE STAR-LINER TELESCOPES

		Standard	De Luxe
6"	Student Model .....	f/8 \$229.50	
6"	with Star-Liner Mount f/8	\$328.50	\$ 450.00
8"	" " " " f/7	\$425.00	\$ 595.00
10"	" " " " f/6	\$536.00	\$ 765.00
12½"	" " " " f/6.5	\$885.00	\$1195.00

16" (Special orders from \$4,250.00 to \$7,900.00)

Finest optics throughout. All mirrors and diagonals figured to ¼ wave or better.


MANUFACTURED IN U. S. A. — SKY TESTED

AND UNCONDITIONALLY GUARANTEED

(Phone LO 3-8429)

## OBSERVING WITH A CLINOMETER

THE Abney-type clinometer is an inexpensive pocket-size device used by builders and engineers to determine vertical angles. It can readily be employed for measuring the sun's altitude if a very dense filter is added, and therefore I have made some tests of its suitability for approximate determinations of geographical position and local time.



The unit consists of a tube five inches long, with a viewing hole at one end and a horizontal wire at the other. A mirror arrangement brings the image of a spirit-level bubble into the same field of view, provided the altitude does not exceed 40°. In observing the sun, its disk is bisected by the horizontal wire, and a knob is turned to place the bubble alongside the wire. The vertical angle is read from an arc of one-inch radius, graduated in degrees, with the aid of a vernier indicating

10 minutes of arc. In the experiments, each observation of the sun consisted of the average of five settings taken in rapid succession.

From a comparison of 55 observed altitudes with computed values, the index correction was found to change from -4' at 8° to +3' at 35°. The mean error of a single observation was  $\pm 5'.2$ . A series of 39 observations of the sun's alti-

A clinometer such as this can be used for a variety of simple experiments in practical astronomy. The observer sights through a small peephole in the right end of the square tube. The inclined gray tube (partly hidden) holds a spirit level which provides an artificial horizon.

tude around noon gave the geographical latitude of my observing station at Weston, Massachusetts, as  $+42^\circ 21'4''$  — only 0'.6 or 0.7 mile south of the true position.

Compared with a bubble sextant, this simple hand-held nontelescopic device yields quite satisfactory accuracy, and is cheaper and much more compact. It has, however, two serious disadvantages. Altitudes greater than 40° cannot be measured, and lack of artificial illumina-

tion for the bubble restricts observations to daytime or bright twilight. Hence, for astronomical applications, only the sun, moon, and Venus are available.

J. A.

## SUNSPOT NUMBERS


The following American sunspot numbers for April have been derived by Dr. Sarah J. Hill, Whitin Observatory, Wellesley College, from AAVSO Solar Division observations.

April 1, 77; 2, 58; 3, 71; 4, 75; 5, 82; 6, 81; 7, 61; 8, 51; 9, 44; 10, 34; 11, 40; 12, 41; 13, 36; 14, 49; 15, 39; 16, 58; 17, 68; 18, 70; 19, 70; 20, 60; 21, 43; 22, 38; 23, 27; 24, 43; 25, 43; 26, 54; 27, 70; 28, 85; 29, 77; 30, 86. Mean for April, 57.7.

Below are provisional mean relative sunspot numbers for May by Dr. M. Waldmeier, director of Zurich Observatory, from observations there and at its stations in Locarno and Arosa.

May 1, 102; 2, 84; 3, 72; 4, 55; 5, 42; 6, 36; 7, 31; 8, 28; 9, 44; 10, 46; 11, 45; 12, 56; 13, 52; 14, 46; 15, 38; 16, 31; 17, 23; 18, 44; 19, 47; 20, 59; 21, 58; 22, 66; 23, 74; 24, 78; 25, 72; 26, 47; 27, 41; 28, 38; 29, 36; 30, 41; 31, 24. Mean for May, 50.2.

Sunspot activity continues to decline and now is back at the level of late 1955. Dr. Waldmeier has made the following predictions of smoothed monthly sunspot numbers for the coming months: July, 59; August, 56; September, 54; October, 52; November, 49.



## Breathtaking Performance —

with an

## ASTROLA REFLECTING TELESCOPE

American Made — Available on Time Payments

Truly breathtaking performance is yours with an ASTROLA reflector. If you are an ASTROLA owner, you enjoy many thrilling views — the finer markings on Jupiter's cloud belts, gaps in Saturn's ring system, delicate details on Mars, such as the canals — all far beyond the grasp of smaller or less perfect telescopes. If you haven't ordered your ASTROLA, why wait? Countless observing thrills will be yours throughout the rest of 1961 and for many years to come — with an all-precision ASTROLA reflecting telescope. For solid observing comfort, nothing matches the De Luxe ASTROLA models, with electric clock drive, large solid-brass setting circles, and smoothly rotating tube, which always allows the eyepiece to be horizontal.

Here we present our new 8" f/6 De Luxe ASTROLA on a portable equatorial mounting with clock drive, setting circles, 8 x 52-mm. finder, and three of the finest orthoscopic oculars. This telescope can be completely disassembled in less than three minutes.

Price, complete, \$590.00

### PORTABLE ASTROLA REFLECTORS

6" Student, f/7 to f/9..	\$194.50
Standard	
6".....	\$325.00
8".....	\$390.00
10".....	\$495.00
12½" Transportable.....	\$1150.00

### NEW PARABOLIC MIRRORS AND DIAGONALS OF PYREX-BRAND GLASS

6".....	\$40.00
8".....	\$92.50
10".....	\$160.00
12½".....	\$250.00

Send for our new 1961 complete color catalogue and full price sheet. Visit our display room when you are in the Los Angeles area. Always in stock are many small reflectors, refractors, books, and accessories.

## CAVE OPTICAL COMPANY

4137 E. Anaheim St., Long Beach 4, Calif.

Phone: GENEVA 4-2613

## DEEP-SKY WONDERS

TO MANY OBSERVERS, globular clusters are the most majestic objects in the sky. In the northern heavens they appear mostly alone — riding like solitary galleons on a vast sea. But the region of Scorpius, which is well up above the southern horizon this month, abounds in both open and globular clusters.

The picture here, from E. E. Barnard's *A Photographic Atlas of Selected Regions of the Milky Way*, gives a hint of the riches of this area. In the upper (northern) part of the photograph is Rho Ophiuchi and its surrounding nebulosity; Antares is at bottom left. To the right of Antares is the rich globular M4, while above and to the right of the star is another globular, NGC 6144, forming a squat triangle with Antares and M4. A third globular is M80, near the right edge of the upper part of the picture.



This photograph of the vicinity of Rho Ophiuchi and Antares was made in 1905 by E. E. Barnard using a 10-inch lens of 50-inch focus, the exposure being  $4\frac{1}{2}$  hours. The nebulosities cannot be seen visually. Reproduced courtesy of the Carnegie Institution of Washington.

M4 (NGC 6121) is a good object for beginning a tour of this field. Easily found by working from Antares, it is at right ascension  $16^h 20^m.6$ , declination  $-26^\circ 24'$  (1950 co-ordinates). Helen S. Hogg lists only six other globulars having diameters greater than its  $22'.8$  (about two-thirds the apparent size of the moon!). But of these six, only M13 is in the northern sky, the others being far to the south.

Mrs. Hogg lists M4's photographic total magnitude as 7.41, and others have recorded visual magnitudes as bright as 6.8. I have seen this object easily with the unaided eye from Kansas, Colorado, and Arizona, and would estimate it as 6.5. In northerly England, T. W. Webb could see it only as "large and dim." But for even small telescopes in the United States it is a splendid subject; in southern latitudes it rivals M13.

The fairly conspicuous NGC 6144 is absent from Norton's *Star Atlas*, although it is only slightly fainter and smaller than M72, the faintest globular in Messier's list. With a magnitude of about 10 and a diameter of  $6'$ , it is quite easy. Look for it at  $16^h 24^m.2$ ,  $-25^\circ 56'$ .

M80 (NGC 6093), at  $16^h 14^m.1$ ,  $-22^\circ 52'$ , is  $5'$  in diameter and said to be of the 8th magnitude, though it appears brighter to me. One meets it easily in sweeping, and it seems to have a concentrated central glow that stands out. It was in M80 that nova T Scorpii blazed in 1860, the only known case of a nova associated with a globular cluster.

There are too many globulars to the east and south of this region to list here, but the sky watcher enjoying an exceptionally clear night should try for NGC 6541, down near the end of the Scorpion's tail. For observers at latitude  $40^\circ$  north, this cluster can climb only about  $6^\circ$  above the horizon. Located at  $18^h 04^m.4$ ,  $-43^\circ 44'$ , it will be a severe test. Yet this globular has nearly the same diameter and magnitude as M13. I would appreciate a card from anyone who can obtain a clear view of this unfamiliar object.

WALTER SCOTT HOUSTON

36 Lawn Ave.  
Middletown, Conn.

## Impossible?

1. M6 Wide-Field Elbow Telescope —  $8^\circ 45'$  field;  $2\frac{1}{16}"$  eye lens; 4 filters;  $15\frac{7}{8}" \times 6\frac{3}{4}" \times 7\frac{3}{4}"$ , 10 lbs.; coated optics; excellent condition ..... \$27.50
2. GSAP (AN-N6) Aerial Gun Camera — 16-mm., magazine loading; 24-volt operation; 35-mm.-f.l.  $f/3.5$  lens. New ..... \$14.00  
Good Condition ..... \$ 9.50
3. Kodak Aerostigmat Lens —  $12" f.l.$ ,  $f/5.0$ ; new, in carrying case with cone, diaphragm ..... \$19.00
4. Kodak Aero-Ektar or Bausch and Lomb Aero Tessar —  $24" f.l.$ ,  $f/6.0$ ; new, in carrying case with diaphragm, cone ..... \$22.50  
Same, but with coated lenses .... \$29.00
5. M3 Panoramic Telescope — coated, new ..... \$ 6.00
6. M76C Telescope — coated, excellent condition ..... \$16.00
7. M17 8x Elbow Telescope — coated, 4 filters, excellent condition ..... \$ 9.50
8. 7 x 50 Binocular Eyepiece — coated, new; by Universal Camera \$ 2.50
9. Bausch and Lomb 7 x 50 Binocular Prism Shelf — 2 right-angle prisms, new ..... \$ 2.50
10. Bausch and Lomb Metrogon Wide-Angle Lens —  $6" f.l.$ ,  $f/6.3$ , in cone with Fairchild high-speed shutter; excellent condition ..... \$49.00

Many unlisted optical, electronic, electro-mechanical items. Showroom open 11:00 a.m.-3:30 p.m., Monday through Saturday.

**THERMOELECTRIC DEVICES INC.**  
**SURPLUS DIVISION**

302 Massachusetts Ave., Cambridge 39, Mass.  
Phone: UN 4-8644 or BE 2-7426





# UNITRON

## MONTHLY REPORT TO OBSERVERS

SEE  
THE BACK COVER  
FOR MORE NEWS  
ABOUT  
UNITRON



### constellation of the month

## SCORPIUS

On a clear, dark July evening, the star-spangled Milky Way forms a luminous arch across the eastern sky. This broad band of light is especially rich in Scorpius, low in the south. The brilliant red star Antares marks the heart of the Scorpion, whose large constellation figure can be traced by the naked-eye watcher with little strain on his imagination.

This fine constellation is a special favorite of many telescope users. It is a memorable experience to roam through its rich star fields and many clusters with even a small refractor, using a large-field eyepiece.

A 3-inch UNITRON will give a rewarding view of the famous globular cluster Messier 80, situated about midway between Antares and 2nd-magnitude Beta Scorpii. This softly glowing ball of light is actually a system of myriads of closely packed faint stars, some 34,000 light-years distant.

Open clusters abound in Scorpius. Messier 7 is especially notable as a rich swarm of points of light. Because this cluster is so far south, your best view may be from a country hilltop with a free southern horizon. For such an expedition, the compact and easily transported 2.4-inch UNITRON is particularly well-suited. Nearby, and also in the Scorpion's tail, is another rewarding open cluster, Messier 6.

Double stars are numerous in this constellation. Even in your finder, the wide pair of 4th-magnitude stars charted as Omega<sup>1</sup> and Omega<sup>2</sup> is an eyecatcher. Not far to its northeast is Nu Scorpii, shown by a 1.6-inch UNITRON as a 41-second double, magnitudes 4 and 6. But a 3-inch in good seeing will add to the latter an 8th-magnitude star two seconds away, and a 6-inch may reveal the brightest star as a one-second pair.

Beta Scorpii is a very unequal double, with a dim 10th-magnitude secondary 14 seconds distant. Fairly high magnification on a 3-inch UNITRON should split it nicely.

Antares itself is a famous but difficult double star, its 6½-magnitude companion a scant three seconds from the dazzlingly bright primary. It calls for a 6-inch UNITRON and a night when the air is very steady.

### HAS IT OCCURRED TO YOU?

There is much to recommend a UNITRON Refractor as the logical choice for the amateur astronomer. A UNITRON, optically speaking, duplicates the performance of larger telescopes of other types. With its long focal length, higher magnifications of planetary and lunar images are obtained with low-power eyepieces. Moreover, there are no mirrored surfaces to become oxidized, no components which require periodic alignment, no secondary optics to cause diffraction patterns, and no folding of the light back on itself through turbulent air with consequent loss of definition. No wonder that you see more and see better with a UNITRON — the telescope that has withstood the test of time.

### Some Popular UNITRON Models

1.6" ALTAZIMUTH (\$7.50 Down)	\$75
with eyepieces for 78x, 56x, 39x	
2.4" ALTAZIMUTH (\$12.50 Down)	\$125
with eyepieces for 100x, 72x, 50x, 35x	
2.4" EQUATORIAL (\$22.50 Down)	\$225
with eyepieces for 129x, 100x, 72x, 50x, 35x	
3" ALTAZIMUTH (\$26.50 Down)	\$265
with eyepieces for 171x, 131x, 96x, 67x, 48x	
3" EQUATORIAL (\$43.50 Down)	\$435
with eyepieces for 200x, 131x, 96x, 67x, 48x	
4" ALTAZIMUTH (\$46.50 Down) with	\$465
eyepieces for 250x, 214x, 167x, 120x, 83x, 60x	
4" EQUATORIAL with clock drive	\$985
(\$98.50 Down), Model 160V, eyepieces as above	
4" PHOTO-EQUATORIAL with clock drive and	\$1175
Astro-camera (\$117.50 Down), with eyepieces	
for 250x, 214x, 167x, 120x, 83x, 60x, 38x, 25x	



### UNITRON accessory of the month


## DUETRON

Here is one of UNITRON's most versatile accessories.

With the handy DUETRON Double Eyepiece, two observers can use the telescope simultaneously and with equal comfort. It is the ideal accessory for those who bought their UNITRON in partnership or for father-and-son observing teams. Makes "star parties" more fun and is the perfect tool for instruction to the beginner.

The two eyepiece holders of the DUETRON may be turned independently to provide convenient positions for both observers.

Priced at \$23.50, this fine UNITRON accessory makes a perfect gift for the amateur astronomer. For double viewing pleasure, you can't beat a DUETRON by UNITRON.



The DUETRON Double Eyepiece is shown in action mounted on a UNITRON 4-inch Photo-Equatorial. A team of four acrobats could be accommodated by adding a DUETRON to the 2.4-inch Guide Telescope seen in the illustration to the right of the main 4-inch refractor. Conceivably, DUETRONS could be connected to DUETRONS to provide for still larger groups, with the 1.6-inch finder used in case the viewers were odd in number. Enthusiasts employing the telescope in this manner should be versed in the art of suspended animation and, for the best results, the entire group should be equatorially mounted, the same as the telescope itself. Should this prove inconvenient, one possible solution is to buy more UNITRONS — a step which would meet with our enthusiastic approval.

# THE MOONS OF JUPITER

Once, not too long ago, there was a king with but two interests in life: astronomy and his three daughters, whom he named after the moons of Jupiter — I, II, and IV. His son III he had banished years before for using one of the royal telescopes as a burning glass to light firecrackers.

Time passed, and the king grew old and almost blind. His study of the sun and stars gave him a perpetual squint, and earned from his subjects the fond nickname of King Leer. Realizing that his observing days were nearly over, he decided to bequeath to his daughters his two most prized possessions — refractors of most marvelous and rare quality (not quite UNITRONS, but still the best in all the land).

"Now have I but two refractors and three daughters," he thought to himself, "and I myself shall want to observe from time to time, when the fog of the years lifts momentarily from my eyes. I know what I shall do; I shall leave the telescopes to the two daughters who love me most. They will let me observe when I like."

And so he called his daughters before him. "How much dost thou love me, dearest I?" he asked. "My love burns brighter than a thousand suns," answered I, whose heart was in reality harder than the bearings in a UNITRON mounting.

"Thou dost please me much," King Leer said. "And now II, how much dost thou love thy father?" "My love," said II with a shrewd glance at I, "burns brighter than a thousand suns seen through a thousand UNITRONS," and the king did not know her heart was harder even than the glass in UNITRON optics.

"Dost thou love me more than this?" asked the king of IV, his favorite. "What has thou to say?" "Nothing," said IV.

"Come, come," said King Leer, "nothing will come of nothing."

"My love is richer than my tongue," replied IV; and this was true, for her heart was as docile as the slow-motion controls of a UNITRON.

The king was enraged, and on the spot he gave his two telescopes (the best in the land, though not up to UNITRON's standard) to his two deceptive daughters, and to IV he left only \$12.50 plus a small pittance every month, along with an outdated Julian Day Calendar, which UNITRON prints annually



for the AAVSO. IV, dejected, left his royal sight.

Next night he went to visit his daughter I, planning to study the star clusters in Scorpius.

"Go away!" she snapped. "I am searching for Venus' moon, and I want no doddering old fool at my eyepiece!"

The king, thunderstruck, left I, without even pointing out that Venus' moon had not been seen since 1764. "I will go to II," he said, brightening. "She will surely let me use her telescope."

But a servant met him at II's door, and said that his mistress was too busy trying to find the missing Messier objects to be bothered with him. The king tried to force entrance to his beloved daughter, but was brought up short by her trained mastiffs. "The little dogs and all," he said, "Tray, Blanche, and Sweetheart, see, they bark at me."

Filled with remorse and repentance, he searched far and wide for IV, realizing now that she alone really loved him. One day, while admiring from afar the observatory atop a 17-story town house, he was astonished to see IV appear at the door. "How came you here?" he cried as he rushed into her arms.

"It was all your priceless gift. But for the Julian Day Calendar, I would never have heard of UNITRON. But for hearing of UNITRON, I would never have sent for their free Observer's Guide and Catalog. But for

the Catalog, I would never have found out about their Easy Payment Plan. I put down the \$12.50 you left me and agreed to pay the small pittance every month. In return UNITRON sent their wonderful 2.4" Altazimuth Refractor, and that was how I met Harry."

"I don't understand," said the king.


"I was observing the star clusters in Scorpius one night with my new UNITRON," she replied. "Harry happened to come by and asked if he could look too. I said it was difficult for two to look through one eyepiece. 'Not with a DUETRON!' he said, and bought me one. We found observing with the DUETRON so pleasant that one day he asked me to become his satellite. He loves astronomy."

"And is that immense observatory on the roof for your UNITRON 2.4" refractor?" asked the king, somewhat puzzled.

"Well, no," replied IV. "It seems that Harry already had a UNITRON 6" Photo-Equatorial with clock drive. We put the DUETRON on that, and we've been happily UNITRONing ever since. Would you care to join us? You can have my 2.4" UNITRON."

The king was delighted, and immediately rushed to the roof to try his new possession. "What magnificent definition!" he cried as he gazed with wonder into the eyepiece. It turned out that he was not going blind after all; the prized refractors he had given his daughters simply did not show the heavens as clearly as a UNITRON.

## ANOTHER NEW UNITRON EXTRA:



In this case, UNITRON's brand new Achromatic Amplifier — a two-element, color-corrected Barlow-type negative amplifying lens designed especially for UNITRON Refractors.

What's good about it is this: When used with a UNITRON eyepiece, it doubles normal magnification to increase the usefulness of each eyepiece. Two types of mounting cells are available, one to fit the UNI-HEX Rotary Eyepiece Selector, and one for the UNITRON Star Diagonal. It's easy to use, and there's never any adjustment of eyepiece (or star diagonal) necessary. What's best about the new Achromatic Amplifier is that it is now included with every UNITRON at no additional cost.

## IT HAD TO HAPPEN: A UNITRON WITH YOUR NAME ON IT!



We've gone and done it. We've personalized the telescope business. For a limited time, when you buy a UNITRON, you can have your name embossed in a plastic name plate attached to the gleaming black barrel at no charge.

People tell us they like it. It's in quiet good taste, yet it seems to give that extra fillip to the pride of ownership people feel for their UNITRONS. To get your name plate simply add this statement to your order: Please attach a name plate to my telescope. This is the name: (print name clearly).

## HOW TO ORDER A UNITRON

Send check or money order in full or use our Easy Payment Plan. Shipments made express collect. Send 20% deposit for C.O.D. shipment. UNITRON instruments are fully guaranteed for quality, workmanship, and performance, and must meet with your approval or your money back. Prices of UNITRON refractors include basic accessories, eyepieces, tripod and mounting, fitted cabinets, and instructions.

## USE OUR EASY PAYMENT PLAN

UNITRON's popular Easy Payment Plan is a convenient and economical way to buy your UNITRON Refractor. The down payment required is 10%. The balance due is payable over a 12-month period, and there is a 6% carrying charge on the unpaid balance. Your first payment is not due until 30 days after you receive the instrument, and if you should want to pay the entire balance due at that time, the carrying charge is cancelled.

## Get UNITRON's FREE Observer's Guide and Catalog

### Contents include:

- Observing the sun, moon, planets and wonders of the sky
- Constellation map
- Hints for observers
- Glossary of telescope terms
- How to choose a telescope
- Amateur clubs and research programs



## UNITRON

INSTRUMENT COMPANY • TELESCOPE SALES DIV.  
66 NEEDHAM ST., NEWTON HIGHLANDS 61, MASS.


Please rush to me, free of charge, UNITRON'S new Observer's Guide and Telescope Catalog 20-R.

NAME \_\_\_\_\_

STREET \_\_\_\_\_

CITY \_\_\_\_\_

STATE \_\_\_\_\_



## Astronomy means observation

Identification of celestial objects during summer evenings is easy with our Sky Publications.

Particularly suited for amateurs and students with small telescopes who desire sky charts in book form is Norton's Star Atlas, a familiar and famous reference handbook. Covering the entire heavens, it shows over 9,000 stars to magnitude 6-1/3, nebulae, and clusters. Descriptive lists of 500 objects and sun, moon, and planet data aid the observer. 108 pages, \$5.25.

For a more detailed atlas, the 16 charts developed by Antonin Becvar and his associates at the Skalnaté Pleso Observatory in Czechoslovakia are available. 35,000 stars to magnitude 7.75, with double, multiple, and variable stars, novae, clusters, globulars, and planetaries, bright and dark nebulae, galaxies, the Milky Way and constellation boundaries, are printed in many colors in the *De Luxe Atlas of the Heavens*, the 16 1/2" x 23" maps being bound in a heavy cloth cover. \$9.75.

Observers at their telescopes will find the inexpensive Field Edition of the Skalnaté Pleso Atlas a most convenient star reference. 16 sheets of stiff paper, 18" x 12 1/4", are printed in white on a black background, so they may be illuminated with a flashlight without spoiling the viewer's dark adaptation. The charts are unbound, \$4.00 per set; two sets for \$7.50.

Antonin Becvar's *Atlas Coeli-II, Katalog 1950.0* is the most complete check list of celestial objects ever offered to the amateur observer. Given, with descriptive data, are the 6,362 stars brighter than magnitude 6.26, with their right ascensions and declinations, magnitudes, and much other useful information; extensive lists of star clusters, nebulae, and galaxies. Cloth bound, 367 pages, \$8.75. The catalogue makes a fine companion to the *De Luxe Atlas of the Heavens*, and they sell together for \$17.50.

Lunar observing is more informative with Elger's Map of the Moon, a large canvas-mounted chart identifying all important lunar features. Notes by H. P. Wilkins on 146 areas are given. The lunar chart itself is approximately 18" in diameter. \$3.00.

In two colors, and over 10" in diameter, the SKY AND TELESCOPE Lunar Map identifies most important features on the moon, including 326 mountains, seas, and craters. The map is good for use with groups at star parties: individual copy, 25 cents; 3 to 100 maps, 20 cents each; over 100, 10 cents each plus postage.

The constellations are easily identified from any location in latitudes 52° to 32° north with the aid of Philips' Planisphere. This world-famous star finder is printed in dark blue and gold, and can be set quickly for any time of year. Directions for using the device are given in four languages. \$3.00.

Decorative as well as informative is the Color Map of the Northern Heavens, a large wall chart showing the northern sky in polar projection to declination -43°. Such important star groupings as Gould's belt, the Scorpion-Centaurus association, the blue stars of Orion, are all indicated by the star symbols, colored vividly to show spectral class. 30" x 34 1/2", only \$1.00.

Write for catalogue of all Sky Publications. All items, except where noted, sent postpaid. Please enclose your check or money order.

## Announcing

## AN ATLAS OF THE MOON'S FAR SIDE

The Lunik III Reconnaissance

On October 4, 1959, the Soviet Union successfully launched its third space probe, Lunik III, into orbit around the moon. Carrying a battery of intricate scientific equipment, including photographic and television systems, it passed near the moon, oriented so as to view a portion of the lunar surface that is perpetually hidden from the earth. On command it photographed the moon's far side for 40 minutes, and these images were later telecast to the earth.

The results of this achievement were compiled and edited for the U.S.S.R. Academy of Sciences by three Soviet astronomers, N. N. Barabashov, Kharkov University Observatory, A. A. Mikhailov, Pulkovo Observatory, and Yu. N. Lipsky, Sternberg Astronomical Institute. Dr. Lipsky summarized this work in SKY AND TELESCOPE for March, 1961, page 133. Now the complete Russian *Atlas of the Opposite Side of the Moon*, translated into English by Richard B. Rodman of Harvard Observatory, is being published jointly by Interscience Publishers, Inc., and Sky Publishing Corporation.

This English edition contains every fact, every illustration, in the Russian original. It includes an introduction, "The Photographs and Their Transmission," "Interpretive Techniques," "The Photometric Cross Sections," "Reduction of the Material," and replicas of all 20 full-page plates with their 30 halftone pictures of hitherto invisible lunar features. These pictures have been obtained by means of electronic filtration processes from the best original negatives taken by Lunik III. A major part of the book is the detailed catalogue identifying and describing 498 lunar formations. The definitive map is given in two forms — both as four full pages in the book, and as a separate 17" x 24" folding sheet. 200 pages (8 3/8" x 11 1/2"), 20 plates. **\$7.00**

### Making Your Own Telescope

Amateurs thinking of constructing a 6-inch reflecting telescope are reminded of Allyn J. Thompson's *Making Your Own Telescope*. Complete step-by-step directions for making and mounting the instrument at low cost are given in easy-to-understand chapters. No complicated mathematics is involved, and no prior knowledge of optics is needed. 211 pages, 104 illustrations (7th printing). **\$4.00**

**SKY PUBLISHING CORPORATION**

Harvard Observatory, Cambridge 38, Massachusetts



# BOOKS AND THE SKY

## PLANETS, STARS, AND GALAXIES

Stuart J. Inglis. John Wiley and Sons, Inc., New York, 1961. 474 pages. \$6.50.

ACCORDING to the jacket, this handsome book is written for nonspecialist readers who "like to speculate for themselves about the universe." That is, the author sets out to acquaint the reader with the major, unsolved problems of astronomy — the frontiers of the subject — without requiring of him any specialized preparation in mathematics or physics.

It is often argued that this is an impossible task. "The frontiers of a science are so far removed from the layman's level of discourse," say the specialists, "that many years of technical preparation are necessary just to comprehend the problems." Such a defeatist attitude is responsible, no doubt, for the ever-expanding undergraduate requirements for science majors, and for the gulf between science and art so eloquently described by C. P. Snow. Surely there is an urgent need in our society to provide good literature that informs the public of just what scientists are doing.

Mr. Inglis is to be commended for the goal he set for himself. He is well qualified, as a college instructor in astronomy and a planetarium lecturer, to undertake the task, and has drawn wisely on the considerable resources of West Coast observatories for illustrations and for the most recent professional thinking in research astronomy.

To some extent, *Planets, Stars, and Galaxies* achieves its goal. The illustrations are of good quality, the diagrams are clear, and there are but half a dozen simple equations in the whole book. The reader needs only an understanding of graphs, ratios, and powers of 10.

The text is clear enough, with few unexplained terms and symbols, but is arranged somewhat too strictly in textbook style, without notable inspiration or elegance. The author uses a strongly deductive approach, proceeding "logically from paragraph to paragraph and from chapter to chapter, so that the discussion of each new topic represents a coherent development of the preceding material," as the statement on the jacket so aptly puts it. Many teachers of elementary astronomy will welcome this coherence, but it immediately structures the subject in a form to be learned rather than enjoyed. In this regard the book is no better than several other recent introductory texts, although it does cover more of the interesting problems of modern astronomy.

What is really needed for the nonspecialist reader, and what Mr. Inglis has not provided, is a contribution to the literature of science, a book that reveals the connected nature of the subject without being pedantic, a book that draws on the excitement of exploration and discovery

from cover to cover, and emphasizes the process of scientific inquiry as well as the results.

After a brief historical introduction, Chapter 2 turns quite logically to basic tools and methods of astronomy, from optical telescopes and Schmidt cameras to spectrographs, photometers, and radio telescopes. Then, in a mere 11 pages, Chapter 3 treats the atom, atomic structure, energy levels and series in the hydrogen spectrum, and nuclear reactions — the proton-proton reaction and (oddly) the last step only of the Bethe carbon cycle. Chapter 4 is a routine description of the physical characteristics of the planets, and is followed by "The Planetary Motions," which starts with Ptolemaic theory and gives a page each to Copernicus, Galileo, and Kepler (without much chance of showing the reader the radical change in astronomical concepts along this historical sequence). This section ends with a summary of the earth's motions (rotation, revolution, solar motion, galactic rotation, and so forth), thus making a start on the basic question of absolute motion treated in the last chapter.

Under the title "Satellites, Natural and Man Made," the next chapter discusses the moon in considerable detail (including R. Baldwin's analysis of crater sizes and N. A. Kozyrev's spectrum of gases in the crater Alphonsus), eclipses, tides, and the known satellites of other planets. The general treatment of artificial earth satellites does not refer to specific launchings, but explains stable orbits and "weightlessness" well, and describes the Van Allen belts as an example of what new data can be obtained.

The foregoing suffices to show the general scheme followed in the remaining 10 chapters, on asteroids, meteors and comets, the sun, stellar spectra, "deviant stars" (variables and novae), multiple stars (binaries and clusters), interstellar material, and galaxies, including the Milky Way.

The publisher's promise of encouraging "intelligent speculative thinking" is limited primarily to three chapters: "The Age and Origin of the Solar System," "The Evolution of Stars," and "The Universe and Relativity." In the first of these, the theories of Buffon, Kant, Laplace, von Weizsäcker, Kuiper, and Hoyle are discussed, together with the solar system features to be explained, and the author partly succeeds in showing how a sequence of theories may slowly approach some ultimate truth, stimulating a great deal of useful research along the way.


The chapter on stellar evolution, one of the best written, starts with modern stellar models and brings in most of the supporting evidence (cluster H-R diagrams, Beta Canis Majoris stars, T Tauri stars, Herbig-Haro objects, and white dwarfs), indicating the wide variety of

Star Atlases and Books on Astronomy	
New: <b>CHANGING VIEWS OF THE UNIVERSE</b> , by C. A. Ronan.....	\$3.95
New: <b>GALAXIES</b> , by H. Shapley.....	\$5.00
<b>TOOLS OF THE ASTRONOMER</b> , by G. Miczaika and W. Sinton.....	\$7.75
<b>SOURCE BOOK IN ASTRONOMY</b> , ed. by H. Shapley.....	
Before 1900.....	\$3.00
1900-1950.....	\$10.00
<b>AMATEUR ASTRONOMER'S HANDBOOK</b> , by J. B. Sidgwick.....	\$12.75
Available again: <b>OBSERVATIONAL ASTRONOMY FOR AMATEURS</b> , by J. B. Sidgwick.....	\$10.75
<b>SKY OBSERVERS GUIDE</b> .....	\$2.95
<b>SUNDIALS</b> , by R. N. and M. Mayall.....	\$3.95
<b>BAA Handbook for 1961</b> .....	\$2.00
<b>AMATEUR TELESCOPE MAKING</b> , Book 1.....	\$5.00
Book 2.....	\$6.00
Book 3.....	\$7.00
<b>Norton's STAR ATLAS</b> .....	\$5.25
<b>Beyer-Graff STAR ATLAS</b> .....	\$15.00
<b>BONNER DURCHMUSTERUNG</b> .....	\$100.00
Write for free list of astronomical literature, including books on telescope making, optics, optical glassworking, pocket books, and paperbacks.	
<b>HERBERT A. LUFT</b>	
67-11 229th St., Oakland Gardens 64, N. Y.	

## OUTER SPACE PHOTOGRAPHY FOR THE AMATEUR

by Dr. Henry E. Paul

In **OUTER SPACE PHOTOGRAPHY FOR THE AMATEUR**, the well-known amateur astronomer, Dr. Henry E. Paul, presents a straightforward guide to sky-picture taking with a camera and simple equipment. Lavishly illustrated. Published by AMPHOTO.



On sale at camera and book stores, or order direct from the publisher.

**AMPHOTO** 33 W. 60th St., New York 23, N. Y.

## CLASSIC AND RECENT CLOTHBOUND BOOKS

Keep in step with happenings in astronomy and the space sciences! Note these new and old, technical and popular, books:

<b>CHANGING VIEWS OF THE UNIVERSE</b> — C. Ronan. 206 pages.	\$4.00
<b>THE WORLD ABOUT US</b> — Sir Graham Sutton, editor. 122 pages.	\$3.95
<b>ASTRONOMICAL PHOTOGRAPHY AT THE TELESCOPE</b> — T. Rackham. 232 pages.	\$7.95
<b>ELEMENTS OF MATHEMATICAL ASTRONOMY</b> — M. Davidson. 225 pages.	\$5.00
<b>THE EXPLORATION OF SPACE</b> — Robert Jastrow, editor. 160 pages.	\$5.50
<b>ADVANCES IN THE ASTRONAUTICAL SCIENCES, Vol. 6</b> — H. Jacobs and E. Burgess, editors. 898 pages.	\$22.50
<b>ROCKET PROPELLANT HANDBOOK</b> — B. Kit and D. S. Evered. 384 pages.	\$12.50
<b>AIRCRAFT AND MISSILE DESIGN AND MAINTENANCE HANDBOOK</b> — C. A. Overbey. 346 pages.	\$9.75
<b>MATERIALS FOR ROCKETS AND MISSILES</b> — R. G. Frank and W. F. Zimmerman. 124 pages.	\$4.75
<b>HANDBOOK OF GEOPHYSICS</b> — U. S. Air Force. 656 pages.	\$15.00
<b>HELICOPTER DYNAMICS AND AERODYNAMICS</b> — P. R. Payne. 422 pages.	\$17.00
<b>AERODYNAMICS OF THE HELICOPTER</b> — A. Gessow and G. C. Meyers, Jr. 343 pages.	\$9.00
<b>THE HELICOPTER</b> — J. Shapiro. 269 pages.	\$4.50
<b>GLIDING: A HANDBOOK ON SOARING FLIGHT</b> — D. Piggott. 261 pages.	\$5.00
An Old-Timer Is Back in a New Edition:	
<b>GALAXIES</b> — H. Shapley. 186 pages.	\$5.00
<b>SOURCE BOOK IN ASTRONOMY 1900-1950</b> — Many writers, edited by Harlow Shapley. 423 pages.	\$10.00
<b>TOOLS OF THE ASTRONOMER</b> — G. R. Miczaika and W. M. Sinton. Replaces <i>Telescopes and Accessories</i> . 294 pages.	\$7.75
<b>GALILEO IN CHINA</b> — Pasquale M. D'Elia, S.J. Translated and revised by M. Sciascia and R. Suter. 115 pages.	\$4.00

Circulars still available. Drop us a card!

## ASTRONOMY CHARTED

33 Winfield St., Worcester 10, Mass., U.S.A.  
Phone: PL 5-6992

the  
review of  
**POPULAR  
ASTRONOMY**



**The Monthly Evening Sky Map** — in its revised and expanded format — set the world of amateur astronomy talking last year. Now — in 1961 — the name has been changed to reflect more clearly its broadened coverage. Here are a few reasons why 5,000 amateurs are enjoying each issue:

- ★ Articles by prominent professional and amateur astronomers.
- ★ Morning and evening sky maps . . . detailed equatorial charts showing planets, Messier objects, variables, stars to 5th magnitude.
- ★ "Through the Three-Inch" . . . informal discussion of constellations and objects of interest.
- ★ Features on telescope making, asteroids, comets, meteors, occultations, variables, and satellites by such staff members or regular contributors as Armand Spitz, Walter S. Houston, Tom Cave, Dr. I. M. Levitt, Leon Campbell, Jr., and David Meisel.

**TRIAL OFFER AND BONUS!**

Send order now, receive current issue free — subscription will start with the following issue. Money returned if not pleased with bonus issue.

One Year: \$3.00      Two Years: \$5.00  
Published Bimonthly      Sample Copy 50¢

**The Review of Popular Astronomy**  
**SKY MAP PUBLICATIONS, INC.**  
P. O. Box 231-S, St. Louis 5, Mo.

observational data to be integrated into one satisfactory theory.

The last chapter, on the universe and relativity, begins with the distribution of galaxies (C. D. Shane's counts); conventionally reviews distance criteria and Hubble's law; discusses the Michelson-Morley experiment, special relativity, the geometry of curved space, the rival theories of cosmology and nucleogenesis of G. Gamow and F. Hoyle; and ends with the question of life elsewhere in the universe. I doubt that this chapter will start much speculation, intelligent or otherwise; the author has pulled so many theoretical rabbits out of his hat up to this point that the reader wouldn't dare question the magic — he simply waits for more. It is never made clear what the ground rules of cosmological speculation are; the emphasis is entirely on explication of conclusions now considered "right" or, in daring moments, "possibly right."

The value of *Planets, Stars, and Galaxies* is considerably enhanced by a glossary of new terms, a list of further readings, many of them articles in *SKY AND TELESCOPE* and *Scientific American* as late as mid-1959, and four or five questions or exercises for each chapter. It also has a good 11-page index, star charts, and (on the inside of the dust jacket) a photographic map of the moon.

THORNTON PAGE  
Wesleyan University

**CAPTURED STARS**

Heinz Letsch. Gustav Fischer Verlag, Vil-  
lengang 2, Jena, East Germany, 1959.  
183 pages. DM 16.00.

**THIS** impressively titled book presents a limited, subjective analysis of the early-model Zeiss planetarium instrument now being manufactured in East Germany. It is a translation by Harry Spitzbardt of the German work *Das Zeiss Planetarium*.

Copyrighted in 1959 by Carl Zeiss, Jena, *Captured Stars* impresses this reader as being a lesser imitation of a more comprehensive, complete book about the same subject, *From the Aratus Globe to the Zeiss Planetarium*. The latter was written by Helmut Werner in close collaboration with the inventor of the Zeiss instrument, the late Walther Bauersfeld. It was copyrighted in 1953, then revised and enlarged in 1957 by Carl Zeiss, Oberkochen, West Germany.

Whether by oversight or intent, *Captured Stars* is limited because it does not contain, among other important elements, a description of the newest and improved Zeiss planetarium, installed in London about two years before the book was printed. Another of these modernized instruments has since been set up at the American Museum-Hayden Planetarium in New York, and its integral-component innovations adapted to two of the other five Zeiss projectors in this country (Chapel Hill and Chicago) and to several planetariums abroad. To be objective and complete, any authoritative text on the planetarium printed after 1957 should contain a description of these advanced features, which significantly increase the scientific utility of the instrument and reproduce the beauty of the night sky far more realistically.

*Captured Stars* is offered as a "considerably condensed text." This is certainly accurate, because some of the important aspects of the planetarium story have been condensed away entirely. The stated purpose of the book is to "arouse an appreciation of the ingenuity of the invention and the cultural achievement of the Planetarium." But so compressed is the text that the name and the work of the ingenious inventor have been omitted completely. Also omitted are descriptions of the diverse cultural potentials of planetariums.

The text reads as if Mr. Letsch had accepted the assignment to analyze scientifically the effects produced by the planetarium, to study available references, and to compile a book on the subject, without having personally experienced the intricacies of the physical operations.

The translation from the German is perhaps too literal, because in many instances the text requires even the initiated person to pause and ponder the accurate but uncommon phrase and sentence structure.

Although the table of contents indicates

With this fascinating, precision-built globe combination you can follow the exciting events of an entire day in the heavens. You read time by observing the earth's relation to the sun. There are two globes, a 6-inch terrestrial and a 12-inch celestial. The earth globe is rotated by an electric motor once every 24 hours. The sun's noontime position is shown for any point on the earth's surface, and because the movable ecliptic ring is connected to the twilight ring, sunrise and sunset and the duration of twilight are shown for all places.

Only two daily settings are necessary — simply move the sun indicator eastward to the next day's date on the ecliptic circle, and turn the earth globe one degree to correspond to the actual rotation of the earth (about 361° in 24 hours).

The movable ecliptic (sun) ring is also marked for the moon's phase positions and the maximum greatest elongations of Mercury and Venus.

Model ST-12-E.C. — 12-inch diameter . . . \$97.50

Write for free brochure.

**FARQUHAR TRANSPARENT GLOBES**

3724 Irving St., Philadelphia 4, Pa.  
Telephone: EVergreen 2-2835

**FARQUHAR EARTH CLOCK**



The latest new

**Spitz** MODEL A-3-P

**PLANETARIUMS**

have been installed at


Amherst College  
Amherst, Massachusetts

Louisiana State University  
Baton Rouge, Louisiana

Temple University  
Philadelphia, Pennsylvania



**Spitz Laboratories**  
YORKLYN, DELAWARE  
Phone: CEder 9-5212



completeness of coverage, the author has ignored the human-interest features of the planetarium. Therefore he fails to satisfy the general reader who is awed, instructed, entertained, and inspired by what the instrument can do, but seeks only a superficial understanding of the mechanical operation of this complex scientific tool. On the other hand, Mr. Letsch has not produced a book adequate for the technician who needs to understand the functional aspects of the mechanism, nor for the professional who wants a complete, authoritative reference work.

Although *Captured Stars*, with 62 pages of illustrations and 86 of text, is an impressive book, it does not accomplish its stated purpose as effectively as its competitor, *From the Aratus Globe to the Zeiss Planetarium*.

A. F. JENZANO  
Morehead Planetarium

#### RADIO ASTRONOMY

**F. Graham Smith.** Penguin Books, Baltimore, 1960. 265 pages. \$1.65, paper bound.

USE of the radio region of the spectrum to complement and extend man's knowledge of galaxies, stars, and the interstellar medium is only about three decades old. It is still a mushrooming field, producing several important discoveries each year. F. Graham Smith, a member of the radio astronomy group at the University of Cambridge, has made a number of important contributions to the subject, especially in determining the positions and flux densities of radio sources and in connection with the solar corona, the earth's ionosphere, and galactic radiation.

In *Radio Astronomy* Dr. Smith attempts to cover in approximately 250 pages every aspect of this very broad subject. He gives the reader an insight into the direction that radio astronomy is taking as well as where it has been. When one considers the breadth of coverage, the length of the book, and the intended audience, the author has done a good job. Any intelligent layman who enjoys reading about astronomy at the SKY AND TELESCOPE or *Scientific American* summary-article level should enjoy this treatment.

It is impossible to write a book on radio astronomy that will be completely up to date at the time of publication, but Dr. Smith comes admirably near this goal. He treats such timely topics as the zodiacal light and the boundary between the earth's upper atmosphere and the outer solar corona, whistlers, Faraday rotation, and radar astronomy. The "belt" or "spur" of continuous radiation that appears in the galactic plane just to the east of the galactic center and extends to high latitudes (a phenomenon that has still not been fully explained), the recent emphasis on measuring angular diameters of radio sources, and the novel technique of aperture synthesis are discussed.

At one point the author speculates on the level of extragalactic radio astronomy knowledge attained by a hypothetical observer on a planet somewhere inside Cygnus A, and on the limits which the noise level of his environment would place on sensitive radio reception.

It is rather unfortunate that the term "radio star" has been used throughout more than half the book; only on page 156 does the author introduce the name "radio nebula." There are a number of misprints and typographical errors, which include reference to the 250-inch, instead of the 250-foot, Jodrell Bank steerable radio telescope (page 115). On page 141, the author names Michigan State University as constructing a single reflector with a collecting area of five acres, for work around 70 centimeters. Michigan State is not so engaged. Perhaps the author means Ohio State or the University of Illinois, but the description does not exactly fit either institution. The amount of atmospheric refraction not only depends "on the air pressure, at ground level, and on the altitude of the star being observed," (page 207) but is also temperature dependent.

From statements made on pages 16 and 178, the reader could easily get the wrong impression that the Fraunhofer lines are formed mainly in the solar corona.

It should be mentioned that the centimeter-wave observations of Saturn and the Helix and Dumbbell nebulae, reviewed on page 182, have not stood the test of time. The diameter of each of the California Institute of Technology paraboloids mentioned on page 252 is 90 feet, not 80.

These few criticisms are minor, however, and F. Graham Smith is to be congratulated for his stimulating treatment of an exciting new field of science.

WILLIAM E. HOWARD  
University of Michigan Observatory

#### RIVAL THEORIES OF COSMOLOGY

**H. Bondi, W. B. Bonnor, R. A. Lyttleton, and G. J. Whitrow.** Oxford University Press, New York, 1960. 64 pages. \$2.25.

THIS little book grew out of a series of lectures given for the British Broadcasting Corp. by William Bonnor, Hermann Bondi, and R. A. Lyttleton, followed later by a discussion among the three of them, moderated by G. J. Whitrow. The rivalry referred to in the title is between the conventional cosmology based on general relativity, supported by Bonnor, and the steady-state theory, adhered to by Bondi and Lyttleton. The publisher recommends the book to readers who have no previous specialized knowledge, but some acquaintance with the basic notions of cosmology is highly desirable, even though no mathematics is needed.

Actually the text is a debate between Bonnor and Bondi. Whitrow acts as an



★ years of EXPERIENCE

★ ENGINEERING know-how

★ basic PATENTS and


★ MANUFACTURING facilities to meet your

## DOMES PLANETARIUMS TELESCOPE

needs for stimulating new  
interest in the  
**ASTRO-SCIENCES**

Call or write : ★  
for descriptive data  
covering a decade  
of design

U.S.  
REPRESENTATIVE  
FOR GOTO  
OPTICAL  
MFG. CO.  
OF TOKYO





impartial moderator, while Lyttleton's contribution is limited to a bizarre piece of speculation about the charges of protons and electrons. The first two men argue about cosmology, but their real disagreement goes much deeper. They represent fundamentally different approaches to scientific theory.

Both men agree on two basic points. First, the primary test of a theory is that it agree with the observed facts. Second, in the absence of clear-cut observational tests, preference is to be given to the simplest theory that fits the available data. The more artificial assumptions a theory has to call upon, the less valuable it is as an explanation. To decide whether an assumption is natural or artificial is not so easy, however, and here Bonnor and Bondi differ strongly.

## Astronomy Films

16-mm. sound, 400-foot reels

I THE SUN; II THE MOON;  
III SOLAR SYSTEM; IV MILKY WAY;  
V EXTERIOR GALAXIES.

**2 x 2 SLIDES**

**35-mm. STRIPS OF SLIDES**

THROUGH 200-INCH AND  
OTHER GREAT TELESCOPES

Catalogues on request.

**International  
Screen Organization**

1445 18th Ave. North, St. Petersburg 4, Fla.

Now six times a year

## Spaceflight

*A bimonthly magazine  
on man's greatest adventure!*

- SPACEFLIGHT is now being published bimonthly — six times a year (January, March, May, July, September, and November). This popular yet authoritative magazine on rockets, astronautics, and space-travel astronomy is written especially for the layman and is edited by members of the British Interplanetary Society.
- Leading authorities provide a comprehensive coverage of all fields of science that play an important part in this thrilling adventure of mankind. Rocketry, artificial satellites, space medicine, atomic fuels, radar controls, exploration of the planets, are all treated in nontechnical language.
- SPACEFLIGHT is mailed directly to you during the month of issue from the printer in Great Britain. Subscription in North and South America: \$3.50, one year; \$6.00, two years. Single copy, 75 cents. (Please specify the issue with which your subscription should start. Back issues available.)

*Sample copy sent on request.*

**SKY PUBLISHING CORPORATION**

49 Bay State Rd., Cambridge 38, Mass.

The problem is that the laws of physics and astronomy are based on observations confined to much smaller distances and shorter time intervals than those involved in cosmology: how then are we to carry these laws over to larger distances and longer times? Bonnor argues that in the absence of information to the contrary, the simplest course is to assume that the conventional laws can be carried over without modification. Since our most general theory of mechanics is general relativity, it is the latter's equations that should be used to investigate the expanding universe.

Bondi, on the other hand, believes that a cosmologist should begin by considering possible models of the universe, and then eliminate those that conflict with observations. From this point of view he believes the simplest possible theory to be that of the steady state, in which the over-all picture of the universe does not change with time. To avoid having expansion thin out the universe, he has to introduce the hypothesis of continuous creation of matter. Bondi considers this hypothesis just as plausible as any other, since it has never been disproved by observation. Bonnor, however, regards continuous creation as a completely artificial hypothesis designed merely to avoid a changing density in the universe.

The cosmological argument will eventually be settled by observation, but the philosophical argument will not. The advance of science is based on a mixture of the points of view represented by Bonnor and Bondi. The question is really one of balance, and the answer involves individual judgment. What blend of the two approaches is likely to be most fruitful for scientific progress? If the majority of astronomers incline toward Bonnor's more conventional view of cosmology, it is because their judgment tells them that Bondi's view of scientific theory is less likely to lead to fruitful results.

IVAN R. KING

University of Illinois Observatory

## HANDBUCH FÜR STERNFREUNDE

Günter Dietmar Roth, editor. Springer-Verlag, Heidelberger Platz 3, Berlin-Wilmersdorf, West Germany, 1960. 360 pages. DM 48.

**M**ANY suitable English-language books exist for the beginning amateur astronomer who wants to make a first acquaintance with the showpieces of the skies. But there are very few for the advanced amateur who needs a systematic manual describing observing methods and explaining problems he can fruitfully attack.

Probably the best volume of this kind was Joseph Plassmann's German classic of 1922, *Hevelius: Handbook for Friends of Astronomy and Geophysics*. Although now out of date in places, it is a mine of useful techniques and ideas, still well

worth reading. However, copies are hard to find.

A new German book, shorter but in much the same spirit as the old *Hevelius*, is Günter Roth's *Handbook for Amateur Astronomers*. The 13 chapters were written by both professional and nonprofessional astronomers. Throughout, free use is made of elementary mathematics, allowing a much wider range of topics than would a purely descriptive approach.

The first section of some 120 pages is devoted to such matters as instruments and photography, time and astronomical co-ordinates. The coverage is similar in many ways to J. B. Sidgwick's *Amateur Astronomer's Handbook* (1955). However, the English book gives many more practical hints on construction and use of telescopes and accessories, while the more systematic German treatment provides a better insight into principles. It also touches on radio telescopes, omitted by Sidgwick.

Practical observing methods occupy nearly 200 pages. R. Müller has contributed an excellent section on solar work, as has W. Petri on solar eclipses. Worthy of special note are the three chapters on the moon, lunar eclipses, and occultations, which are much superior to the treatment of the same subjects in Sidgwick's *Observational Astronomy for Amateurs*. The planets are dealt with rather briefly, but in excellent perspective. A serious omission is of instructions for observing central-meridian transits of Jovian features and the determination of their rotational periods; this information is provided in detail by Sidgwick. Another noteworthy chapter deals with artificial satellites, but the account of visual meteor work is far less helpful than Sidgwick's.

One strong point of Roth's book is its treatment of variable star observing. Here the Argelander step technique is well explained. The difference in brightness between the variable and each of several comparison stars is estimated in terms of a small arbitrary but constant unit, the step, and a scale of step values can be constructed for the comparison-star sequence. Hence it is possible to determine the light curve of a variable and evaluate times of maximum or minimum, without knowing the magnitudes of any reference stars. Unfortunately the Argelander method is far too little known by American amateurs, although it makes possible much fascinating work outside the routine of organized variable star observing.

Roth's *Handbook* gives a revealing view of the different development of amateur astronomy in Europe and America. For the American with a reading knowledge of German, this volume will prove a fruitful source of new ideas and new techniques. Its value is enhanced by numerous illustrations, extensive bibliographies, and a well-chosen appendix of tabular data.

J. A.

## NEW BOOKS RECEIVED

**ARTIFICIAL SATELLITES**, Michael W. Owen, 1960, Penguin. 128 pages. \$1.25, paper bound.

A British astronomer views the past, present, and future of rocketry and space exploration. Numerous diagrams and photographs illustrate the popularly written text.

**THE PHYSICAL PRINCIPLES OF ASTRONAUTICS**, Arthur I. Berman, 1961, Wiley. 350 pages. \$9.25.

In textbook format, this thorough exposition of the basic principles of astronautics is for students and general readers with a background in introductory physics and calculus. Particular attention is given to the foundations of mechanics, orbit theory, and propulsion dynamics.

**GALAXIES**, Harlow Shapley, 1961, Harvard University Press. 186 pages. \$5.00.

Written by a former director of Harvard Observatory, this is one of the early titles in the Harvard Books on Astronomy, published in 1943. It has now been revised extensively.

**MAN'S VIEW OF THE UNIVERSE**, R. A. Lytle, 1961, Atlantic-Little, Brown. 108 pages. \$3.95.

Illustrated with a familiar selection of celestial photographs and diagrams, this brief summary brings the general reader an up-to-date, authoritative picture of astronomical knowledge and cosmological theories.

**SURFACE OF THE MOON**, V. A. Firsoff, 1961, Hutchinson and Co. Ltd., 178 Great Portland St., London W.1, England. 128 pages. 21s.

A well-known British student of the moon discusses the lunar surface from a geological viewpoint, with special reference to the tectonic grid system. The bibliography has 102 entries.

**THE IMPACT OF THE NEW PHYSICS**, Frank Hinman, 1961, Philosophical Library. 174 pages. \$4.50.

A California School of Medicine biologist, using atomic physics, cosmology, and geology, theorizes on the origin of life.

## SKY-GAZERS EXCHANGE

Classified advertising costs 30 cents a word, including address; minimum charge, \$4.00 per ad. Only one for sale ad per issue for each advertiser. Remittance must accompany order. Insertion is guaranteed only on copy received by the 20th of the second month before publication; otherwise, insertion will be made in next issue. We cannot acknowledge classified ad orders. Sky Publishing Corporation assumes no responsibility for statements made in classified ads, nor for the quality of merchandise advertised. Write Ad Dept., Sky and Telescope, 49 Bay State Rd., Cambridge 38, Mass.

**FOR SALE:** Witherspoon 6-inch refractor objective. Air-spaced doublet in brass cell. Focal length 102 3/4 inches. Like-new condition. \$150.00. Louis W. Breck, M.D., 1220 N. Stanton, El Paso, Tex.

**CONSIDERING ASTRONOMY** as a career? *Vocational and Professional Monographs: Astronomy*, by Dr. Freeman D. Miller, describes personal qualifications, scholastic training, and job opportunities. \$1.00 postpaid. Send to Box B, Sky and Telescope, 49 Bay State Rd., Cambridge 38, Mass.

**OPTICAL COMPONENTS** of amazingly close tolerances. Astro-Research Laboratory, 4601 70th St., La Mesa, Calif.

**FIBERGLASS** telescope tubes, finest available. W. R. Parks, 18015 S. St. Andrews Place, Torrance, Calif.

**FOR SALE:** Sky and Telescope, 1952-57, except October, 1954. Best offer. L. G. Pardue, 641 Falcon, Miami Springs, Fla.

**10" REFLECTOR** for sale: f/8, fully rotating tube. All castings aluminum and bronze; worm gears bronze, ratio 1 to 60. Ball and thrust bearings, cone-type clutch. Setting circles, slow motion. If interested, send for photograph or call for demonstration. Robert G. Fisher, 148 S. Second St., Coplay, Pa.

**AN ELEMENTARY SURVEY OF CELESTIAL MECHANICS**, Y. Ryabov, 1961, Dover. 164 pages. \$1.25, paper bound.

This account of motions in the solar system is intended for the reader who wants something more than the brief statements in popular books, but not a technical monograph. Using only simple mathematics, it should be understandable to informed amateurs. The author explains the basic ideas of gravitation, planetary orbits and their perturbations, and describes in some detail the motions of artificial satellites. This is an English translation of a Russian book first published in 1959.

**TRAITÉ DE MÉCANIQUE CÉLESTE**, F. Tisserand, 1960, Gauthier-Villars, 55 Quai des Grands-Augustins, Paris 6, France. Vol. I, 474 pages, 50 NF; Vol. II, 547 pages, 60 NF; paper bound.

This is a new edition of the *Treatise on Celestial Mechanics*, which since its appearance in 1888-90 has been a classic advanced textbook on motions in the solar system. Volume I, after an introductory hundred pages on undisturbed orbital motion, deals with the problem of three bodies; the second volume is concerned with the shapes and rotations of celestial bodies. The text is in French.

**EL TELESCOPIO**, Miguel Valdez, 1960, Miguel Valdez, P.O. Box 1325, Antofagasta, Chile. 162 pages. \$1.75, paper bound.

Directions in Spanish are given for the construction of an amateur's reflecting telescope, using this small mimeographed book.

**STELLAR ATMOSPHERES**, Jesse L. Greenstein, editor, 1960, University of Chicago Press. 724 pages. \$17.50.

In this advanced textbook, chapters by 20 astronomers discuss in detail the interpretation of stellar spectra, with such related topics as magnetic fields and rotation of stars. While labeled Vol. VI, this is the second part to appear of the nine-volume *Stars and Stellar Systems*, edited by G. P. Kuiper and Barbara M. Middlehurst.

**FRENCH REFRACTOR:** 4 1/4-inch altazimuth; slow motions; massive oak tripod; geared elevating pillar maintains eyepiece at observer's own height. 60x to 300x eyepieces, finder, counterweight. Perfect definition. \$325.00. Small telescope also available. Stephen Smith, Box 1656, Delray Beach, Fla.

**BOOKS WANTED!** Attention astronomical observatories, professionals, amateurs: Collector of rare astronomy works desires to purchase your duplicates or trade disposable volumes. Write, describing fully. Barchas Library, Rte. 4-216, Tucson, Ariz.

**REFRACTOR BARGAIN** — 60-mm. altazimuth, 910-mm. f.l., 150x, only \$59.00 postpaid. Peninsula Scientific, 2421 El Camino Real, Palo Alto, Calif.

**PARABOLIC**, elliptical, spherical mirrors, refiguring. Dual-slow-motion equatorial mounts, Aluminum castings. John McQuaid, 2003 High, Logansport, Ind.

**DALL-KIRKHAM** Cassegrainian, f/15: 6" clear aperture, \$145.00; 8" c.a., \$210.00; 10" c.a., \$290.00; 12" c.a., \$405.00. Maksutov f/23, designed by John F. Gregory: 6" c.a., \$525.50; 8" c.a., \$775.00; 10.8" c.a., \$1,200.00. 25% down, a year to pay the balance. Manufactured by Three "B" Optical Co., Box 508, Mars, Pa.

**ALUMINUM TUBING:** Eighteen sizes, 1" through 10". PESCO-A, Box 1363, Ann Arbor, Mich.

**CELESTAR-6** by Fecker. Catadioptric system, 6" aperture, 90" focal length, 35" tube. No spider diffraction. Electric drive, circles. 3 eyepieces, turret. \$450.00 f.o.b. Melvin Mattox, 3801 Atlantic, Long Beach, Calif.

**MOUNTED** first-quality 6" telescope objectives, \$350.00. Correspondence invited. 24 years experience. Earl Witherspoon, Sumter, S. C.

**LIBRARIES, BOOKSELLERS**, save time. Order Target: Earth (Kelly-Dachille) directly from Target: Earth, Box 355, Carlsbad, Calif.

## Attention!

Introducing a complete line of **UNITRON** telescopes, parts, and accessories

You will be pleased to discover a special line of UNITRON telescopes and many extra accessories, now on display at your new UNITRON dealer:

**Polaris Telescope Shop (Main Store)**  
14319 Michigan Ave.

Dearborn, Mich. (Phone: LU 2-8466)

**NOTE:** A new direct mail order **Supply Center** has been added to aid astronomical groups and amateur telescope makers select supplies, parts, and accessories. For details write: **Polaris Telescope Shop #2**, P. O. Box 2364, Dearborn, Mich.

## Precision Diagonals

You will get the best possible performance from your telescope with one of our clear **FUSED QUARTZ** diagonals. Accuracy guaranteed 1/20 wave.

Ellipse 1.25" x 1.77" . . . \$12.00  
Ellipse 1.5" x 2.12" . . . \$15.00

**PYREX**-brand glass diagonals, 1/8-wave accuracy.

Ellipse 1.25" x 1.77" . . . \$ 6.00  
Ellipse 1.5" x 2.12" . . . \$ 9.00

Without aluminum coating, deduct \$1.00.

Send for our complete list of supplies, quartz mirrors, blanks, oculars, coatings, and accessories.

**E & W OPTICAL CO.**

2420 East Hennepin Ave.  
Minneapolis 13, Minnesota

Don't fail to investigate the new 1961 Magnusson telescopes, mountings, and clock drives.

## IMAGINE THESE FEATURES ON LOW-PRICED REFLECTORS

- ★ Mountings made with 1 1/2" shafts.
- ★ Cast-aluminum holding straps machined to permit easy tube rotation.
- ★ Clock drives with two slow motions, one for scanning the sky and another for photographic work.
- ★ Drives with hardened and ground worm threads that you would expect to find only in instruments costing twice as much.
- ★ Drives with large motors that will accommodate telescopes up to 12 1/2".

All of these extras are made possible without increasing the prices because of our new tools and production methods.

**ALSO AVAILABLE:** Mirror kits from 6" to 12 1/2", with the new channeled glass lap for fast grinding without sticking. Each kit contains a spherometer for measuring depth of curve. The graduated knob with screw can be used on a Foucault tester.

Write for prices — parts sold separately.

## SETTING CIRCLES

Made of choice aluminum or brass, machined and polished all over. Hour circles, machine-scribed with hour, half-hour, and five-minute marks. Declination circles scribed in degrees 0-90-0-90. Numbers stamped with 1/8" dies. Holes reamed standard sizes. Fastened with setscrews. State hole sizes.

	Aluminum	Brass
5" circles, set of two	\$15.00	\$19.50
6" circles, set of two	\$17.00	\$22.10
8" circles, set of two	\$24.00	\$31.20

**O. MAGNUSSON**

14570 West 52nd Ave., Arvada, Colo.



### 3-inch Astronomical Reflector

60 to 180 Power  
An Unusual Buy!

Assembled — ready to use! See Saturn's rings, the planet Mars, huge craters on the moon, star clusters, moons of Jupiter, double stars, nebulae, and galaxies! Equatorial-type mounting with locks on both axes. Aluminized and over-coated 3"-diameter f/10 primary mirror, ventilated cell. Telescope comes equipped with a 60x eyepiece and a mounted Barlow lens, giving you 60 to 180 power. A finder telescope, always so essential, included. Sturdy, hardwood, portable tripod.

FREE with Scope: Valuable **STAR CHART** plus 272-page "HANDBOOK OF THE HEAVENS" plus the book "HOW TO USE YOUR TELESCOPE."

Stock #85,050-Y.....\$29.95 ppd.



### INFLATABLE CELESTIAL GLOBE

Useful for the student beginning to study the stars, and the newly interested amateur. 12" heavy-gauge vinyl globe shows major constellations, with stars to 4th magnitude.

Stock #60,166-Y.....\$1.40 ppd.

### 7 x 50 BINOCULARS — TREMENDOUS BUY! War-Surplus American-Made



Excellent for deep-sky and Milky-Way observation. Brand new! Crystal-clear viewing — 7 power. Coated optics. An excellent night glass — recommended for satellite viewing. Individual eye focus. Exit pupil 7 mm. Approx. field at 1,000 yds., 376 ft. Carrying case included. American 7 x 50's usually are \$195.

Stock #1533-Y..(Tax included).....Only \$55.00 ppd.

6 x 30 Binoculars — similar to above and a bargain!  
Stock #963-Y..(Tax included).....\$33.00 ppd.

### NOW! A FULL SELECTION OF ORTHOSCOPIC EYEPieces


Wide, flat field — better correction under high magnification — excellent eye relief.

The orthoscopic eyepiece is one of the most important and best corrected eyepieces for astronomical work. These are of a four-element design, with coated lenses, and are standard 1 1/4" outer diameter, precision made of chrome-plated brass and aluminum.



Stock #30,364-Y.....	4 mm.....	\$14.50 ppd.
Stock #30,404-Y.....	6 mm.....	14.50 ppd.
Stock #30,405-Y.....	12.5 mm.....	14.50 ppd.
Stock #30,406-Y.....	18 mm.....	14.50 ppd.
Stock #30,407-Y.....	25 mm.....	14.50 ppd.

### Rack-&-Pinion Eyepiece Mounts



Real rack-and-pinion focusing with variable tension adjustment; tube accommodates standard 1 1/4" eyepieces and accessory equipment. Lightweight aluminum body casting (not cast iron); focusing tube and rack of chrome-plated brass; body finished in black wrinkle paint. No. 50,077-Y is for reflecting telescopes, has focus travel of over 2", and is made to fit any diameter or type tubing by attaching through small holes in the base. Nos. 50,103-Y and 50,108-Y are for refractors and have focus travel of over 4". Will fit our 2 1/2" I.D. and our 3 3/8" I.D. aluminum tubes, respectively.

#### For Reflectors

Stock #50,077-Y..(less diagonal holder)....\$8.50 ppd.  
Stock #60,049-Y..(diagonal holder only).... 1.00 ppd.

#### For Refractors

Stock #50,103-Y..(for 2 1/2" I.D. tubing)....\$12.95 ppd.  
Stock #50,108-Y..(for 3 3/8" I.D. tubing).... 13.95 ppd.

### NEW LOW PRICE FLASHLIGHT POINTER

... Point It Out With Projected Arrow



Ideal for pointing to interesting features on movie and slide projection screens. Excellent lecture tool. For teacher use on maps, etc. Flashlight focuses an arrow where you point it.

Stock #60,117-Y.....\$5.95 ppd.

### BARGAIN-PRICED CAMERA SHUTTER



For only \$5.95 you get a shutter worth much more. Speeds: Bulb, 1/25, 1/75, and 1/200th second. Iris diaphragm, 2 to 16 mm. Front threaded opening, 22-mm. diameter; rear threaded flange, 25-mm. diameter. Supplied with retaining ring.

Stock #70,423-Y.....\$5.95 ppd.

### SPECIAL SALE! CRAMER MOTORS Make Your Own CLOCK DRIVE — ONLY \$4.00



Excellent value for the serious amateur. With this motor you can make your own clock drive for about 1/4 the cost of a purchased one. Type 117-P Cramer motor (without clutch), left rotation, 1/15 r.p.m., 115 volts, 60 cycles, 2.7 watts, with a 1/2" shaft plus an extra 1 1/4" long shaft, setscrews, and directions for adapting to your telescope.

Stock #60,205-Y.....\$4.00 ppd.

### Take Pictures Through Your Telescope with the EDMUND CAMERA HOLDER for TELESCOPES



Bracket attaches permanently to your reflecting or refracting telescope. Removable rod with adjustable bracket holds your camera over scope's eyepiece, and you're ready to take exciting pictures of the moon. You can also take terrestrial telephoto shots of distant objects. Opens up new fields of picture taking!



#### SUN PROJECTION SCREEN INCLUDED

White metal screen is easily attached to holder and placed behind eyepiece. Point scope at sun, move screen to focus... and you can see sunspots!

All for the low, low price of \$9.95

Includes brackets, 28 1/4" rod, projection screen, screws, and directions. Aluminum; brackets black crinkle painted.

Stock #70,162-Y.....\$9.95 ppd.

### PRISM STAR DIAGONAL

For comfortable viewing of stars that are high in the sky and overhead. Fits refracting telescopes using standard-size (1 1/4" O.D.) eyepieces, or you can make an adapter for substandard refractors. Contains an excellent high-quality aluminized right-angle prism. The tubes are satin chrome-plated brass. Body is black wrinkle cast aluminum. Optical path length of the system is about 3 1/2".



Stock #70,077-Y.....\$12.00 ppd.

### Operate Your Clock Drive From a 12-Volt Auto Battery



These small inverters supply 110-volt, 60-cycle electricity from your car battery to operate your clock drive anywhere. (Can also be used to operate electric shaver.) 20-watt rating. Size: 2 1/2" x 2 1/2" x 3", 2 lbs.


Stock #50,346-Y... (for 12-volt battery)...\$14.95 ppd.

### NEW SQUARE-SHAPED RETAINING RINGS Set of 20 only \$1.25



Custom-designed square-cross-section retaining rings provide far greater bearing surface than the usual round variety. Made of special carbon steel, these retaining rings really hold your lenses in position. Set includes four each of five different sizes: .618" O.D. x .035" sq.; .878" O.D. x .062" sq.; 1.136" O.D. x .055" sq.; 1.400" O.D. x .062" sq.; and 1.665" O.D. x .071" sq. Complete set provides superior results previously unobtainable, and fits practically all available tube sizes.

Stock #60,207-Y.....\$1.25 ppd.




### THE PLANETARIUM PORTRAYS THE FASCINATING STORY OF EARTH IN ACTION

Here is the story of cosmic motions — how the moon revolves around the earth, and the earth around the sun, with our planet rotating simultaneously. With this instrument, the observer sees a three-dimensional moving demonstration of how seasons, day and night, and moon phases occur. This handsome gear-and-chain-driven unit is supported by a smartly finished wood base. The sun is 6" in diameter, and the earth is 4". Planetarium stands 12" high, is 8" wide; arm is 18" long. A completely illustrated, delightfully informative handbook included.

Stock #70,415-Y.....\$29.95 ppd.

### 6X FINDER TELESCOPE



Has crosshairs for exact locating. You focus by sliding objective mount in and out. Base fits any diameter tube — an important advantage. Has 2 holders, each with 3 centering screws for aligning with main telescope. 20-mm.-diameter objective. Weighs less than 1/2 pound.

Stock #50,121-Y.....\$8.00 ppd.

### TELESCOPE LIGHTING SYSTEM Help Preserve Your Night Vision




Fits any telescope; takes one size D flashlight battery. Case fastens to tripod leg with wing nut and clamp. Two grain-of-wheat lamps and housings with 39" cords. One lamp rheostat controlled. By drilling a small hole in finder tube at the plane of the reticle, you can illuminate the crosshairs, controlling their brightness. Other lamp stays at constant brightness so you can illuminate your setting circles, check eyepieces, examine charts, or make notes.

Stock #50,219-Y.....\$2.95 ppd.

# EDMUND SCIENTIFIC CO.




## THE ORIGINAL GOODWIN RESOLVING POWER BARLOW



For many years, this achromatic coated Barlow lens has been the pride and joy of serious astronomers. Due to the death of the designer, Mr. Frank Goodwin, it has not been available for the past two years. NOW we can offer these lenses, in exact accordance with the original specifications. Remember, this Barlow is achromatic, coated, mounted in a blackened tube, and as optically perfect as only precision craftsmen can make it. Complete with instructions, in a 4"-long adapter tube for standard 1 1/4" eyepieces.

Stock #60,122-Y.....\$23.50 ppd.

## DOUBLE AND TRIPLE YOUR TELESCOPE'S POWER WITH A BARLOW LENS



WHAT IS A BARLOW? A Barlow lens is a negative lens used to increase the power of a telescope without resorting to very short-focal-length eyepieces, and without the need for long, cumbersome telescope tubes. Referring to the diagram above, a Barlow is placed the distance P inside the primary focus of the mirror or objective. The Barlow diverges the beam to a distance Q. This focus is observed with the eyepiece in the usual manner. Thus, a Barlow may be mounted in the same tube that holds the eyepiece, making it very easy to achieve the extra power. The new power of the telescope is not, as you might suppose, due to the extra focal length given the objective by the difference between P and Q. It is defined as the original power of the telescope times the quotient of P divided into Q.



Beautiful chrome mount. We now have our Barlow lens mounted in chrome-plated brass tubing, with special spacer rings that enable you easily to vary the power by sliding split rings out one end and placing them in other end. Comes to you ready to use. Just slide our mounted lens into your 1 1/4" I.D. tubing, then slide your 1 1/4" O.D. eyepieces into our chrome-plated tubing. Barlow lens is nonachromatic.

Stock #30,200-Y...Mounted Barlow lens....\$8.00 ppd.

## BINOCULAR-TO-CAMERA HOLDER

For Exciting Telephoto Shots  
Will fit any camera



Bring distant objects 7 times nearer with a 35-mm. camera, 7 x 50 binocular, and our BINOCULAR-TO-CAMERA HOLDER. Ideal for photographing constellations, star clusters, the moon, as well as cloud formations, wildlife, vistas.

Camera and binocular attach easily. Use any binocular or monocular — any camera, still or movie. Take color or black-and-white. Attractive gray crinkle and bright chrome finish, 10" long. Full directions for making telephoto pictures included.

Stock #70,223-Y.....\$11.50 ppd.

## MORE POWER FROM YOUR JAPANESE TELESCOPE

Mounted Barlow for Japanese Telescopes

By inserting this single-element lens in the eyepiece end of your Japanese telescope, and putting your regular eyepiece in the end of the Barlow tube, you can increase your telescope's power up to three times. Thus, instead of 60x, you will get 120x or 180x. Barlow is mounted in two pieces of telescoping brass tubing each 4" long, satin-chrome plated on the outside. Inner diameter of large tube and outer diameter of small tube are 0.965", the standard size for most Japanese telescopes. Measure yours before ordering. 0.965" is approximately 31/32" or 24.5 mm.

Stock #30,370-Y.....\$6.00 ppd.

## Learn Astronomy The "Fun" Way 53 STAR & PLANET FLASH CARDS



Stock #60,210-Y.....

Complete Set Only \$1.00

Make learning fun and easy. Just look at the picture, then turn the card for identification — full details given on the back. Set includes 29 constellation cards, 10 on the solar system, 10 star cards, and 4 of other celestial objects.

Stock #60,210-Y.....\$1.00 ppd.

## Suppress Internal Reflection FLOCK-PAPER TUBE LINING

Superior To Dead-Black Flat Paint

Special paper with a black velvetlike surface that absorbs light and prevents unwanted reflections and glare. Ideal as a screen on which to project star images for a realistic effect. Easily glued or cemented inside your tube. FEATURED in April, 1961, *Review of Popular Astronomy* (page 13) by G. R. Wright. Comes in sheets, 27" x 36".

Stock #60,068-Y.....per sheet.....\$1.00 ppd.

## STAR TIME CALCULATOR



This handy slide rule automatically makes the conversion from star or sidereal time to standard time. Saves the amateur astronomer the time and trouble of calculations. Size: 10 1/2" x 3 3/4", plastic-coated cardboard.

Stock #40,399-Y.....\$1.50 ppd.

## CLOCK-DRIVEN EQUATORIAL MOUNT ON PEDESTAL BASE IDEAL FOR YOUR 6" OR 8" REFLECTOR

Accurate electric clock drive and heavy-duty mounting. Operates on household current. Follows stars smoothly. Pedestal is 24" high. Polar-axis shaft diameter 1". Setting circles included.

Stock #85,111-Y.....\$74.50 f.o.b. Barrington, N. J.

Same mount and clock drive on 32" hardwood tripod.

Stock #85,081-Y.....\$69.00 f.o.b. Barrington, N. J.

Same mount on metal pedestal, no clock drive.

Stock #85,108-Y.....\$45.00 f.o.b. Barrington, N. J.

Same mount on tripod, no clock drive.

Stock #85,023-Y.....\$39.50 f.o.b. Barrington, N. J.



## ACT NOW! SAVE PLENTY!

Here they go — all first-class aerial camera lenses and astronomy sets. Priced to go, along with many other real buys, discounted 25 per cent and now discontinued.

AERIAL CAMERA LENS, f/6, 24" f.l. — was \$39.50.

Stock #85,059-Y.....While they last.....\$25.00 f.o.b. Utah

ASTRONOMY SET, projects moon travel, eclipses, stellar wonders — was \$5.00.

Stock #70,233-Y.....Now \$3.75 ppd.

DE LUXE ASTRONOMY SET, down from \$10.00.

Stock #70,234-Y.....Now \$7.50 ppd.

GET LISTINGS OF OTHER GREAT SAVINGS IN BARGAIN BULLETIN


## Mounted Ramsden Eyepieces

Standard 1 1/4" Diameter

Our economy model, standard-size (1 1/4" O.D.) eyepiece. We mounted two excellent quality plano-convex lenses in black anodized aluminum barrels instead of chrome-plated brass to save you money. The clear image you get with these will surprise you. Directions for using short-focal-length eyepieces are included with both the 1/4" and 1/2" models.

Stock #30,204-Y.....1/4" focal length.....\$4.75 ppd.

Stock #30,203-Y.....1/2" focal length.....\$4.50 ppd.



## SETTING CIRCLES TO FIT YOUR EDMUND TELESCOPE OR TELESCOPES WITH SIMILAR-SIZED MOUNTS



A — Heavy-Duty



B — Light-Duty

(A) FOR HEAVY-DUTY MOUNT (fits Edmund 6" reflector, 4" refractor). Black plastic. Hour circle has two scales, bottom 0 to 24 hrs., top 0-6, 6-0, etc. Declination circle has numbers every 10°, lines each degree. All filled white. Hole is 1.9"; 3" outer diameter. Two pointers with 3/4" holes.

Stock #60,188-Y.....\$3.00 ppd.

(B) FOR LIGHT-DUTY MOUNT (fits Edmund 4 1/4" reflector, 3" refractor). Black plastic, scales same as above. 1"-diameter hole; 3" outer diameter. Two pointers with 1 1/4"-diameter holes.

Stock #60,187-Y.....\$3.00 ppd.

## NOW! LENS ERECTOR FOR TERRESTRIAL VIEWING WITH YOUR REFLECTOR

This Edmund development adds real convenience to viewing objects on the earth. Just put the lens erector in your eyepiece holder, insert eyepiece, and focus normally. You see everything right side up and correct as to left and right. Made of polished chrome-finished brass telescoping tubing that will fit any standard 1 1/4" eyepiece holder. Tubing is 9 1/2" long and slides 3" into eyepiece holder. Erecting system consists of two coated achromats.

Stock #50,276-Y.....\$9.95 ppd.

## GIANT ERFLE EYEPIECE 1 1/2" F.L.

War-Surplus Bargain — Gov't. Cost Approx. \$100

Large telescopes should have one of these for low-power viewing. Apparent field of view 65°. Also use with our 24"-focal-length Aerial Camera lens to make a 16-power wide-field telescope or a 27-power scope with one of our 40"-focal-length Aerial Camera lenses. Low-reflection-coated, 5-element lens system. Field lens of Eastman Kodak's rare-earth glass for better aberration correction. Has diopter scale. Smooth focusing, 3/4" movement. Outside diameter of attaching threads, 3" — 32 threads per inch. Clear aperture of eye lens 2", field lens 1-25/32". Weight 3 1/2 lbs.

Stock #50,091-Y.....\$9.95 ppd.



## ADAPT GIANT ERFLE TO STANDARD TELESCOPE EYEPIECE SIZE

This adapter lets you use our Giant Erfle with any telescope having a standard 1 1/4"-diam. eyepiece holder.

Stock #50,358-Y.....\$3.95 ppd.

## FREE CATALOG "Y"

144 Pages! Over 1000 Bargains!

Fantastic variety — rarely have so many lenses, prisms, optical instruments, and components been offered from one source. Probably the greatest assembly of bargains in all America. Imported! War surplus! Hundreds of other hard-to-get optical items. Many science and math learning and teaching aids.

Write for Free Catalog "Y".



ORDER BY STOCK NUMBER . . . SEND CHECK OR MONEY ORDER . . . SATISFACTION GUARANTEED!

**BARRINGTON • NEW JERSEY**

# ENGINEERED FOR THE LAST WORD IN BREATHTAKING PERFORMANCE . . . PRICED FOR FIRST CHOICE IN VALUE . . . FULLY EQUIPPED WITH MANY EXTRA FEATURES!

You'll Marvel At How  
The Superb Optics Of  
This Portable RV-6

## 6-inch DYNASCOPE®

Give The Same Exquisite Definition  
As Far More Expensive Instruments!

This new addition to the Dynascope line has won immediate recognition from schools, colleges, and professionals, as an outstanding achievement in a 6-inch telescope. Although it was only recently introduced, our files are already filled with complimentary letters from excited amateurs and professionals all over the country. Each one is truly amazed at the superior optical performance of this new RV-6 6-inch Dynascope! Here is large aperture in a quality instrument at a price that compares with many 4-inch telescopes. *And this low cost includes such exclusive extra features as electric drive (patented), setting circles, and rotating tube!* There are no "extras" to run up your cost!

The superb optical system resolves difficult objects with definition that is absolutely breathtaking. The close tolerances of the precision construction assure an accuracy and smoothness of operation once associated only with the finest custom models. The heavy-duty mount, complete with electric drive, provides the stability so essential for satisfactory viewing, yet there is easy portability because in a matter of minutes the entire telescope can be dismantled into three easy-to-handle sections.

Only Criterion's engineering ingenuity, coupled with volume production and modern manufacturing methods, makes this handsome 6-inch model available at such reasonable cost. You can order it with complete confidence that it will live up to your expectations in every way, for this assurance is *guaranteed* under our full-refund warranty. Send your check or money order today. Or use our liberal time-payment plan and take months to pay.

### YOU COULD PAY \$100 MORE WITHOUT GETTING ALL THESE SUPERIOR FEATURES

(Except on Another Dynascope)

1. EXQUISITE OPTICAL SYSTEM INCLUDES F/8 6-INCH PARABOLIC MIRROR made of PYREX-brand glass, accurate to better than  $1/4$  wave, zircon-quartz coated, and guaranteed to reach all theoretical limits of resolution and definition. Teamed with elliptical diagonal, mounted in 4-vane adjustable spider.
2. NEW DYN-O-MATIC ELECTRIC DRIVE with smooth self-acting clutch that engages and disengages automatically as you seek different objects. Will not interfere when manual operation is desired. Entire drive housed in trim aluminum case. Plugs into ordinary household socket.
3. CHOICE OF ANY 3 EYEPIECES from the following: 70X-18-mm. Achromatic Ramsden; 100X-12.7-mm. Achromatic Ramsden; 140X-9-mm. Achromatic Ramsden; 210X-6-mm. Orthoscopic; 320X-4-mm. Orthoscopic. All are precision-threaded, give sharp images to extreme edges.
4. SOLID NEW EQUATORIAL MOUNT, extra-reinforced design to provide pillarlike stability. No annoying side play or wobble. Adjusts easily to any latitude.
5. SETTING CIRCLES for both right ascension and declination. Handsomely engraved and finished in fine aluminum.
6. 6x30 FINDERSCOPE, ACHROMATIC, COATED, with accurate crosshairs and fine focus. Durable cast-aluminum bracket with 6 adjusting screws allows positive collimation.
7. ROTATING TUBE FOR MAXIMUM VERSATILITY AND VIEWING COMFORT. Solid-cast chrome-finished rings are generous 1" wide with felt lining. Newly designed construction, with over-sized knurled adjusting knobs, affords maximum rigidity and allows for quick disassembly and portability, with or without tube.
8. STRIKINGLY HANDSOME WHITE 50" BAKELITE TUBE with porcelainized Duralite finish, durable yet light. Walls are  $1/8$ " thick, completely insulated and anti-reflective blackened inside.
9. BRASS RACK-AND-PINION EYEPIECE HOLDER has precision-cut rack and gear for easier, smoother, more positive focusing. Takes standard  $1\frac{1}{4}$ " oculars, negative or positive.
10. STRONG, VIBRATION-FREE, ALL-METAL TRIPOD with easily removable legs. Provides sure, steady support, plus lightweight portability.

## CRITERION MANUFACTURING CO.

331 Church St., Hartford 1, Conn.

Manufacturers of Quality Optical Instruments

Including . . . • **ELECTRIC DRIVE (Patented)**  
• **SETTING CIRCLES** • **ROTATING TUBE**

A Complete Instrument, No Costly Accessories Needed!

Model RV-6 Complete  
with Dyn-O-Matic Electric Drive  
and All Features Described Below

**\$194<sup>95</sup>**

f.o.b. Hartford, Conn.  
Shipping Wt. 55 lbs.  
Express Charges Collect  
No Packing or Crating  
Charges

### ENJOY IT NOW FOR ONLY \$74.95 DOWN

No need to put off the thrills of owning this magnificent instrument! Send your check or money order today for only \$74.95 as full down payment . . . pay balance plus small carrying charge in your choice of 6, 12, or even 24 monthly payments. Same unconditional guarantee applies, of course. Or order today by sending your check or money order with coupon below.

Sound too good to be true?  
Then read what these delighted  
DYNASCOPE owners have to say:

"I never dreamed such a brilliant, clear image could be had with a telescope costing under \$200. It is one of these rare bargains that you find only once or twice in a lifetime."

B.S. — New Jersey

"A friend of mine has a 6-inch telescope without electric drive, for which he paid over \$500. He agrees that my new DYNASCOPE not only outperforms it, but is a much better-looking and better-built instrument."

J.L. — New York

"Congratulations on the excellence of your workmanship. The optics are truly amazing. I never expected such wonderful performance."

L.H.N. — Massachusetts

#### Criterion Manufacturing Co.

Dept. STR-24, 331 Church St., Hartford 1, Conn.

☐ Please send me, under your unconditional guarantee, the RV-6 6" Dynascope. Full payment of \$194.95 is enclosed.

☐ I prefer your easy terms! Enclosed is \$74.95 as down payment with understanding that I will pay balance (plus small carrying charge) over 6, 12, 24 months (check choice).

☐ Send FREE ILLUSTRATED LITERATURE describing the RV-6 6" Dynascope and all the telescopes in the Dynascope line.

Name .....

Address .....

City ..... State .....

# GLEANINGS FOR ATM's

CONDUCTED BY ROBERT E. COX

## AN 8-INCH "BEAVERTAIL" DALL-KIRKHAM REFLECTOR

FROM the rich legacy of designs left to amateur astronomers by Russell W. Porter, many have chosen the Springfield mounting, which combines sturdiness with the convenience of having the eyepiece in a fixed position. Usually Newtonian optics are used, introducing the disadvantage that the telescope tube must be supported at its upper end and provided with a long, awkward counterweight.

To overcome this, Porter himself suggested what he called the "beavertail" design (*Amateur Telescope Making — Book II*, page 375), employing Cassegrainian instead of Newtonian optics. The short tube can be supported near the middle, where a diagonal mirror intercepts the light beam to send it out along the declination axis. When it reaches the polar axis, the beam is reflected at a right angle to the eyepiece.

The awkward counterweight is now replaced by one hanging downward from the central mounting assembly, looking like a beaver's tail in Porter's drawing. It is filled with lead to provide the required weight.

This design impressed us as an inter-

esting challenge for our third telescope. The advantage of observing in a comfortable position is indisputable, especially for detailed lunar and planetary work. Hence we felt that this compact instrument would permit us to do more useful observing.

We did not have large metal-working machines, nor the patience to make patterns and castings, so we decided to use the relatively new plasticized wood-particle board (Novaply, Granite Board, Timboard, and the like). It is used extensively where moisture or heat causes natural wood to warp or rot quickly, yet costs less than the best quality plywood. It is flat and uniform, much denser than natural wood, and devoid of grain. Although lacking the ultimate fastener strength of actual wood, it proved satisfactory for the tube and mounting parts.

The movable section of the mounting between the telescope tube and the pier — the heart of the Springfield design — we call the *yoke*. The hollow declination axle, which holds the tube to the yoke, has one radial and one thrust ball-bearing, as well as two silicone-greased 10"



While George Doschek observes at the main eyepiece of this Springfield reflector, his father Antony looks through the finder. The rubber-tired dolly is used to move the telescope to and from its indoor storage place. Leveling is done with the knurled knobs on the base. Photograph by Garry's.

## DE LUXE

### Reflecting Telescope Kits

De luxe kits contain:

- ★ 2 mirror blanks of PYREX-brand glass
- ★ 7 large metal cans of optical abrasives
- ★ Fast-polishing cerium oxide
- ★ Red rouge and pitch

Size	Thickness	Price
4 1/4"	3/4"	\$ 6.00
6"	1"	\$10.50
8"	1 3/4"	\$18.75
10"	1 3/4"	\$33.65
12 1/2"	2 1/8"	\$39.95

ADD POSTAGE: 1st, 2nd, and 3rd postal zones from Detroit, add 10%; 4th, 5th, and 6th, add 15%; 7th and 8th, add 20%; or we will ship postage collect.

Send for free catalog of supplies.

### ASHDOWNE BROS.

18450 Grand River

Detroit 23, Michigan



### SAFEGUARD YOUR EYES!

Don't use your binoculars in glaring sunlight without this **Filtrol**. Unfiltered, uncontrolled, bright sunshine on desert, mountain, or sea will strain your eyes. This Variable

Density Filtrol (Polaroid) varies the amount of light that you need for best viewing without glare. You alone adjust exactly how much light reaches your eyes. Clear aperture, 18 mm. U. S. Gov't. surplus; cost over nine times what we ask. Filtrol to fit over the eyepieces of any binocular, plus instructions. **Only \$1.00 ppd.**

### Jupiter BLUE Filter

A popular, useful accessory for every telescope, especially for viewing the Red Spot on Jupiter. Improves color contrast, reduces glare. Helps you to see detail in the belts of this planet and on Mars. 29.5-mm. diameter, 1 mm. thick. Polished plane parallel. World-famous manufacturer. Unmounted. **Only \$2.00 each, ppd.**

Remit with Order


**HARRY ROSS** Scientific & Lab Apparatus  
61-H Reade St., New York 7, N.Y.

### PARABOLIC MIRRORS OF PYREX-BRAND GLASS

Diam.	Focal Length	Price
6"	48"	\$49.98 postpaid
8"	48"	69.98 postpaid
10"	60"	125.00 postpaid

These mirrors are accurate to 1/4 wave length or better and are free of defects including chips, scratches, and pits. Aluminized reflective coating is protected with quartz overcoating. We invite you to test and compare these with mirrors selling for more. Returnable within 15 days if not pleased.

### EXTRA-THICK PYREX-BRAND GLASS ELLIPTICAL DIAGONALS



Made of select fine annealed material. Flat to 1/4 wave or better and free of any defects. Aluminized and quartz overcoated. Postpaid prices listed below.

### 4-VANE SPIDERS

Simple, sure design. Rigidly constructed. Can be adjusted up or down, tilted or rotated, and locked in place. Postpaid.



Minor Axis	Diagonal	Spider
1 1/4"	\$ 5.95	\$10.95
1 1/2"	8.00	11.50
1 3/4"	10.00	12.95
2"	14.95	14.95
2 1/2"	17.95	16.95
2 3/4"	23.95	22.50

WANTED: Used lenses, eyepieces, objectives, telescopes, books, and the like. Send full descriptions and prices.

Write for free catalog.

### NYE OPTICAL CO.

7310 Broadway, Lemon Grove, Calif.



## OPTICAL FLATS

Pitch Polished, Beral Coated

### Rectangular shape

$1\frac{1}{4}$ " x  $1\frac{1}{4}$ "  
 $\frac{1}{4}$ " wave \$1.75 each  
 $\frac{1}{2}$ " wave \$4.50 each  
 Postpaid.



### Elliptical shape

Heavy edges to minimize temperature effects.

$1\frac{1}{4}$ " minor axis.  
 $\frac{1}{4}$ " wave \$4.00 each  
 $\frac{1}{2}$ " wave \$6.50 each  
 Postpaid.



**BERAL COATINGS** — same optical characteristics as aluminum — mechanically more durable — not over-coated — may be removed without harming glass surface. Prices for Beral coating telescope mirrors: 4" diameter—\$4.50, 6"—\$5.00, 8"—\$6.50, 10"—\$8.50, and 12½"—\$12.50 each, f.o.b. Skokie.

**LEROY M. E. CLAUSING**

8038 Monticello Ave.

Skokie, Ill.

## GET A PERFECT FIGURE ON YOUR MIRROR WITH THIS FINE FOUCAULT TESTER



- Exact zonal measurements
- Machined parts of brass and aluminum

Completely assembled **\$27.50** p.p.d.

**KIT** of all parts **\$14.50** p.p.d.

MIRRORS, MIRROR KITS, BINOCULARS, TELESCOPES, ACCESSORIES

Free Catalogue


**UNIVERSITY OPTICS**

2122 East Delhi Road  
 Ann Arbor, Michigan



Model #6810 mounting for tube assemblies to 10-inch aperture. Cast-aluminum construction, 1.5-inch precision-ground shafting, and 3-inch thrust surfaces render maximum stability in a portable mount.

*T. J. Optical Laboratories*  
 P.O. Box 481  
 18175 Pembroke Circle  
 Huntington Beach, Calif.



The level of concrete in the pier of the Doschek telescope is indicated in this scale drawing. The upper inset shows the secondary mirror's spider support arrangement, the lower one the shape of the pier on its base.

16-gauge polished aluminum plates that act as friction-bearing surfaces. A thin sheet of Teflon between them would be better, but adjustment of the take-up nut against the thrust bearing now provides smooth motion of the tube in declination.

A similar friction-bearing arrangement holds the yoke onto the slanted upper surface of the pier, except that the polar-axis fastener is a solid pin with thrust bearings at each end. The cap end may be fastened to the yoke and the other driven by a clock gear and clutch inside the pier. In our manually controlled version, both polar-axis and declination bearing circles are equipped with outside ring gears turned simply by pinions, whose shafts are held in sleeve bearings through the yoke.

The controls are seen in the photo-

graphs on the right side of the yoke: a handle for movement in declination and a knob and bevel-gear reduction for right ascension. The ring gears were made from 24-pitch straight gear tracks, bent around the 10"-by- $\frac{3}{4}$ " circles and then fastened by long pins driven through holes drilled between the gear teeth every three inches. The small amount of backlash incurred by bending and opening up the teeth of the gear track is not detrimental in view of the frictionally damped motions of the tube and yoke.

The Springfield design requires that the axis-bearing holes be drilled accurately at right angles to the tube and yoke as well as to each other. If a slight error has been made, however, the particle board permits easy correction by sanding the friction-bearing circles. Such an error is detected as a decentering of the several

images of the optical train that are seen by looking down the polar axis without an eyepiece while the tube is rotated in declination.

In the beavertail of the yoke we have 44 pounds of counterweight balancing the weight of the telescope tube and mirror. An easily removable aluminum cassette on the yoke (dark in the pictures) holds the eyepiece focusing rack, the inverting flat, and a slide-out holder for

Prefit all components by tacking them together with long finishing nails; no joint should allow a 0.002" feeler gauge to pass through. Matching joints should be rabbeted to provide greater strength.

After an assembly has been prefitted, the edges of the joints should be sized with white wood glue, or hot glue if available, and allowed to dry, after which the tears and flows should be sanded down. If clearance and pilot holes for the screws


have been drilled during the prefitting, the fastenings can be tightened immediately after gluing and assembly. Use flat-head screws on 2" or 3" centers wherever possible, countersink the heads, and fill the holes with plastic wood.

All holes should be made on a drill press so as to lead the screws properly. Avoid the practice of "towing" the joints, as this causes strains in the assembly. With care and proper centering, the large holes in the tube and yoke that admit the radial and thrust bearings can be bored with an ordinary swing cutter in a drill press. Clamp all parts firmly to the drill table to avoid oversized holes, as the particle board is too soft to hold the pilot drill rigidly on center.

The board's cut edges are porous, requiring a good semi-liquid sealer to be used copiously on all exposed edges after assembly. The sealer is then sanded and painted. Our telescope was given a primer of lacquer sealer, a coat of lacquer surfacer, and a final coat of Multicolor, a tough, leathery finish.

Desiring the best optics available, we asked Daniel E. McGuire, a professional optician, to make the components, the finished system having an equivalent focal length of about 185" and a focal ratio of 24. The field of view with a 26.6-mm. eyepiece is about 15 minutes of arc.

The pyrex primary, 7 $\frac{3}{4}$ " in diameter, is corrected to Dall-Kirkham specifications;



The "yoke" of the telescope, showing the convenient controls for right ascension (knob) and declination (handle). The cassette holding the eyepiece and polar-axis diagonal appears dark above center. The "beavertail" extends downward to the left, holding a number of eyepieces and, at its lower end, a 44-pound counterpoise bolted on.

filters. When this cassette is removed, the focal plane falls almost at the lower edge of the "table" of the yoke, where the image may be examined or photographed.

The square tube is strong and rigid, and is readily fitted with accessories. If ground lights are annoying during observing, the round ventilation holes can be closed with thin sheet-aluminum shields. Note that the lower end of the square tube (a shape easy to construct) is cut on the bias. The part nearest the yoke carries a 26-pound counterweight that is attached to the bottom of the tube after the primary mirror is set in place.

The optics require that the secondary mirror's mounting at the upper end of the tube be extremely sturdy. The spider legs are steel rods  $\frac{3}{8}$ " in diameter threaded into 1" standard pipe sleeves, which are bolted to the corners of the tube (out of the optical path). At the center, these rods hold a Micarta wafer 1" thick to carry the secondary mirror cell and the usual lateral and longitudinal screw adjustments. We wrapped the spider legs closely with coarse black wool yarn, and found that this eliminated most of the visible diffraction pattern. Presumably the diffraction now produces an imperceptible haze dispersed over the field of view.

All of the joined parts of the telescope should be cut to fit as accurately as possible. Even though wood can be "pulled" to fit, this technique should be avoided.

### Read This Advertisement

Here is a combination of a Barlow and a particular ocular which gives outstanding results. It consists of our new Barlow and our 16.3-mm. ( $\frac{3}{4}$ " focal length) Erfle eyepiece. While the Barlow was not specifically designed to work with this eyepiece, it does so to an astonishing degree. All images are sharp and hard to the very edge of the field.

The Barlow gives magnification up to slightly over three times that of the ocular alone. It is achromatic, coated, and mounted to the U. S. standard size of 1.250 inches.

The modified Erfle eyepiece has a field of 75 degrees with excellent eye relief. The combination gives an equivalent focal length of slightly over 6 millimeters. Many users state it is far superior to any shorter focal length ocular of equivalent magnification.

The Barlow sells for \$16.00 postpaid, and the Erfle for \$15.30 postpaid. Both are guaranteed to perform as stated above or money refunded.

#### ORTHOSCOPIC OCULARS—All hard coated, standard 1 $\frac{1}{4}$ -inch outside diameter.

28-mm. ....	\$15.60	10.5-mm. ....	\$16.80	4-mm. ....	\$17.70
16.3-mm. (Erfle) ....	\$15.30	7-mm. ....	\$17.70	Barlow 3x .....	\$16.00

### Telescopes

6565 Romaine Street  
Los Angeles 38, Calif.

Warranted to equal or surpass any oculars obtainable anywhere or money refunded.

Finished mirrors, mirror kits, spiders, elliptical flats, focusing devices, aluminizing.

Send for catalogue.

### DE LUXE SKYSCOPE

As illustrated — with new plastic tube, 6x finder, heavy all-metal tripod, equatorial mounting, and 60x eyepiece.

Price **\$44.75**

Unconditionally guaranteed.  
100% American made.

Write for free brochure showing our standard model which has been sold world-wide for over 20 years. Skyscope still remains outstanding in value by any comparison.

**THE SKYSCOPE CO.,  
INC.**

P. O. Box 55S,  
8104-13th Ave. TE 7-0213  
Brooklyn 28, N. Y.





## OUR LENSES ARE UNEQUALED

So say the many purchasers at universities, laboratories, and government institutions. Prices and quality defy comparison.


# ASTRONOMICAL OBJECTIVES

... Air-Spaced ...

Specifically designed for those who demand an objective lens for genuine astronomical observation.

### WHY BUY FROM US?

On the premise that the objective lens is the heart of the telescope, any prospective builder would certainly expect one that will perform to exacting specifications. Some may ask, "Why a refractor over a reflector telescope?" In the first place, a refractor is much easier to build and cheaper to maintain over the years. A reflector contains many small parts that are subject to wear or loosening, while the refractor is generally more compact, easy to store or carry, and needs little attention. When exploring the possibility of building an astronomical refractor telescope, the question is, "Which objective lens is best?" The answer is simple, and attested to by thousands of satisfied people in the know. "A Jaegers Astronomical Objective" is the only answer. Examine these pertinent facts:



Each lens is thoroughly tested and guaranteed to resolve to Dawes' limit. They are corrected for both the C and F lines (secondary chromatic aberration). The zonal spherical aberration and the chromatic variation of spherical aberration are negligible. The cells are machined to close tolerances so that they fit directly over or into our standard aluminum tubing, eliminating any mounting problems. Test a lens, or have any qualified person test it; we are certain that you will be satisfied. If not, take advantage of our money-back guarantee. We offer the lowest-priced, hand-corrected, precision, American-made astronomical objective, mounted in an aluminum cell. Our reputation for high-quality lenses has established us as the most reliable source in the industry.

### HARD COATED ON 4 SURFACES

3 1/4" APERTURE f/15 — 48" F. L.	Cat. No. S1447 Mounted and Not Coated	\$28.00 ppd.
	Cat. No. S1471 Mounted and Coated	32.00 ppd.
4 1/8" APERTURE f/15 — 62" F. L.	Cat. No. S1472 Mounted and Not Coated	\$60.00 ppd.
	Cat. No. S1473 Mounted and Coated	69.00 ppd.
6" APERTURE f/10 — 60" F. L.	Cat. No. S1861 Unmounted and Coated	\$150.00 ppd.
	Cat. No. S1860 Mounted and Coated	175.00 ppd.
6" APERTURE f/15 — 90" F. L.	Cat. No. S1863 Unmounted and Coated	\$150.00 ppd.
	Cat. No. S1862 Mounted and Coated	175.00 ppd.

## "BIG" ACHROMATIC TELESCOPE OBJECTIVES

We have the largest selection of diameters and focal lengths in the United States available for immediate delivery. Perfect magnesium-fluoride coated and cemented astronomical lenses made of finest crown and flint optical glass. Not mounted. Each lens is guaranteed to resolve to the theoretical limit.

● ALUMINUM TUBING and CELLS are in stock. ●

Cat. No.	Diam.	F.L.	ppd.	Cat. No.	Diam.	F.L.	ppd.	Cat. No.	Diam.	F.L.	ppd.
S1476	2 1/8"	10"	\$12.50	S1520	2 1/8"	50"	\$12.50	S-957	3 1/4"	40"	\$30.00
S1112	2 1/8"	11 3/4"	12.50	S-851	3-1/16"	15"	21.00	S1155	4"	34 1/2"	60.00
S1110	2 1/8"	13"	12.50	S1158	3 1/8"	19 1/2"	28.00	S1460	4 1/4"	36"	60.00
S-958	2 1/8"	15 1/2"	9.75	S-822	3-3/16"	24 1/2"	22.50	S1159	4 3/8"	42"	60.00
S1145	2 1/8"	20"	12.50	S1092	3 1/4"	26"	28.00	S1225	4 3/8"	42"	67.50
S-952	2 1/8"	23 1/2"	12.50	S1093	3 1/4"	28"	28.00	S1474	5-1/16"	24 3/4"	75.00
S1431	2 1/8"	30"	12.50	S1139	3 1/4"	30"	28.00	S1475	5-1/16"	24 3/4"	85.00
S1432	2 1/8"	40"	12.50	S-955	3 1/4"	34 1/2"	28.00				

\*Not coated.

## Astronomical Kits



● **FREE OFFER:** Now you can save many tedious hours of grinding time. Tools and mirror blanks No. S2053 and No. S2054 have a generated f/8 curve. Take advantage of this extra-special offer at no additional cost. For those who desire to grind a curve to their own specifications, the standard blanks are also offered without the generated curves.

**EACH KIT CONTAINS:** Mirror blank, plate-glass tool, eyepiece lenses, first-surface diagonal mirror, assorted abrasives, tempered pitch, inspection magnifying lens.

Cat. No.	Mirror Diam.	Thickness	ppd.
S2053	4 1/4" (f/8 curve)	3/4"	\$ 7.50
S2093	4 1/4" (flat)	3/4"	7.50
S2054	6" (f/8 curve)	1"	11.95
S2094	6" (flat)	1"	11.95
S2055	8" (flat)	1 3/8"	19.50
S2056	10" (flat)	1 3/4"	30.75*
S2057	12" (flat)	2 1/8"	54.75*

\*f.o.b. Lynbrook, N. Y.

## Equatorial Mount & Tripod



### FOR TELESCOPES UP TO 4"

Constructed of cast aluminum and finished in black. The tripod is war-surplus, one of the sturdiest of its size ever built. Made of prime oak with bronze fittings, 3 adjustable locking legs with steel shods. Extended 50", closed 36". Shipping weight is 13 lbs.

Cat. No. S2128 EQUATORIAL MOUNT	\$17.50 ppd.
Cat. No. S2131 TRIPOD	15.00 ppd.

### SAVE MONEY!

Cat. No. S2083 Combination EQUATORIAL MOUNT and TRIPOD	f.o.b. Lynbrook, N. Y., \$27.50
--	---------------------------------

## Reflector Telescope Kits

Excellent quality mirrors, polished, aluminized, silicon-monoxide coated. Each kit contains mirror, diagonal, and lenses for eyepiece. No metal parts. Mounting instructions included.

Cat. No.	Diam.	F.L.	ppd.
S2284	3-3/16"	42"	\$ 9.95
S2285	4 1/4"	45"	14.75
S2286	6"	60"	26.25

## Astronomical Mirrors

These mirrors are of the highest quality. Aluminized with silicon-monoxide protective coating. You will be more than pleased with their performance.

Cat. No.	Diam.	F.L.	ppd.
S1429	3-3/16"	42"	\$ 8.70
S1506	4 1/4"	45"	13.50
S1507	6"	60"	25.00
S2188	6" parabolic	48"	49.50

## Mirror Mounts

Cast aluminum, holds all of our mirrors with metal clips. Completely adjustable, assembled.

Cat. No. S1634 3-3/16"	Mount for 4 1/2" Tubing	\$4.00 ppd.
Cat. No. S1633 4 1/4"	Mount for 5" Tubing	4.00 ppd.
Cat. No. S1632 6"	Mount for 7" Tubing	6.50 ppd.



### ● FREE CATALOG ●

Millions of lenses, prisms, eyepieces, rack-and-pinion eyepiece mounts, cells, aluminum tubing, mirrors, binoculars, telescopes, parts, accessories, and so forth. Write today!

We pay POSTAGE IN U. S. — C.O.D.'s you pay postage. Satisfaction guaranteed or money refunded in 30 days.

**A. JAEGER'S**  
6915 MERRICK RD., LYNBROOK, N. Y.



a  $1\frac{1}{4}$ " perforation at its center is used for mounting the diagonal or coude mirror.\* The primary's focal length is about  $40\frac{1}{4}$ ", f/5.2. The Vicor secondary mirror is  $2\frac{1}{4}$ " in diameter, half an inch thick; it is convex, with a negative focal length of 13". The axial spacing from the primary is 30.625", the amplification factor being about 4.6.

The next two mirrors are the diagonal, 1/20-wave flat Vicor, with a  $2\frac{1}{4}$ " minor axis, placed 27" from the secondary, and the inverting mirror on the polar axis, which is of  $\frac{1}{2}$ -wave pyrex, minor axis  $1\frac{1}{4}$ ". It is located 13" from the coude diagonal, and the final focal plane is about 5" upward along the polar axis. The total length of the optical path is thus only about 76".

It is difficult to describe the beauty and perfection of lunar and planetary details in this telescope. On nights when the seeing is excellent, the crepe ring of Saturn is perceptible at a glance. The red spot on Jupiter is distinctly defined. Because of the long focal ratio, eyepieces show little chromatic aberration, and no color at all is seen on the terminator of the crescent Venus, with only a bare suggestion of color on the planet's edge at magnifications over 300x. Mars shows much detail, and many subtle markings on the moon can be studied.

Needless to say, Messier objects and the like are dimmed by the long focus, but the instrument's stability permits its adaptation to a Newtonian attachment. The central field will be good at f/5.2. In fact, the telescope is so stable and well-damped that twice we have followed the Echo I satellite at 176x by manually guiding the tube at the finder scope while the other observer watched through the Springfield system.

Since September, 1959, our wooden "beavertail," which weighs over 300 pounds, has been hauled from its garage home over rough driveway and lawn the year round; its optical surfaces have been cleansed with soap suds and distilled water twice, yet there has been no observable change in the critical alignment originally made on stars.

ANTONY and GEORGE DOSCHEK  
84B Kingston Ave.  
Pittsburgh 5, Pa.

\*This method of mounting the flat coude mirror through a small hole in the primary has often been recommended by me. It is useful for those who are considering making compound reflectors with unperforated primaries. The hole is small, and centering is not so critical as with a large Cassegrainian perforation. This system gives a very stable diagonal, eliminates a second set of spider vanes, and permits reduction of what otherwise might be too great an effective focal length. It is important, of course, that before fine grinding begins the hole be drilled from the back of the mirror to within  $\frac{1}{8}$ " of the surface, then cut through from the front after figuring is completed.

R. E. C.


## U. S. SURPLUS BARGAINS

IN HIGH-QUALITY OPTICAL PARTS AND EQUIPMENT



**f/5, 12" f.l. AEROSTIGMAT  
COATED LENS  
BY EASTMAN KODAK**

In  $8\frac{1}{2}$ "-long cone,  $11\frac{1}{4}$ " to 6-3/16" diam. No shutter. Iris diaphragm opens  $\frac{1}{2}$ " to  $2\frac{1}{8}$ ". 4-element Tessar design. Front lens clear aperture  $2\frac{3}{8}$ "; rear, 2-5/16". Price **\$30.00**



**WEAVER  
2.2X SCOPE**

Lightweight scope, excellent for big game hunting. Reticle is crosshair type. Can be adjusted for windage and elevation. Universal focus can be used with fixed mount. Barrel diameter  $\frac{3}{4}$ "; length 10 7/8"; weight  $7\frac{1}{2}$  oz. Field of view 23 ft. at 100 yds. Eye relief  $3\frac{1}{4}$ ". Coated lenses. Brand new. Price **\$12.95** postpaid



**f/6.3, 6" f.l. METROGON  
COATED LENS  
BY BAUSCH & LOMB**

In lens cone with iris and shutter. Lens cone O.D.  $11\frac{1}{8}$ "; depth,  $5\frac{3}{8}$ ". Price **\$30.00**



**METAL PARABOLOIDAL MIRROR**


This is made of hastelloy hard metal. 12" diameter.  $5\frac{3}{4}$ " focal length. Manufactured by Bausch & Lomb. Approx. weight,  $3\frac{1}{2}$  lbs. Brand-new condition. Price **\$10.00**

**LENS ASSEMBLY, 48" FOCAL LENGTH, f/6.3**

This was used with the K40 aerial camera, which takes a 9" x 18" plate. Manufactured by Eastman Kodak. Approx. dimensions, 14" diam., 26" long. Contains built-in filters and iris. Lenses mounted in aluminum housing. Approx. weight of unit, 125 lbs. All in original crates. Approx. shipping weight, 200 lbs. Price **\$150.00**

**BELL & HOWELL  
f/8 TELEPHOTO LENS**

Coated lenses. Focal length 36". Completely mounted with iris and shutter. Approx. weight 25 lbs. Excellent condition. Price **\$49.50**



**KODAK AERO-EKTAR f/2.5, 6" f.l.**



Mounted in barrel. Hard-coated lenses, color corrected. Complete with dust caps. Red and yellow filters. Price **\$25.00**

**BAUSCH & LOMB f/6, 24" f.l.  
AERO-TESSAR LENS SET**

The clear diameter of front lens is 4"; that of rear lens is  $3\frac{3}{8}$ ". The set consists of two metal-mounted elements. Price **\$20.00**



**8-POWER WIDE-FIELD ELBOW  
TELESCOPE, COATED LENSES**



Field of view  $8^{\circ} 45'$ . Large focusing Erfle eyepiece. Eye lens diameter 2-1/16" with diopter scale +2 - -4. Four built-in filters—clear, neutral, red, amber. Length  $15\frac{3}{8}$ ". Width  $6\frac{3}{8}$ ". Height  $7\frac{3}{8}$ ". Weight approx.  $9\frac{1}{2}$  lbs. Excellent unused condition. Price **\$39.50**

**NAVY INFRARED RECEIVER,  
TYPE C-3**

Produces visible image from infrared image in complete darkness. For seeing in dark with infrared light source. Nothing to wear out. Operates on two flashlight batteries. Has high-voltage power supply and image tube (1P25) and elaborate optical system. Weighs only  $6\frac{1}{4}$  pounds and is only 11" long, 8" high,  $5\frac{1}{2}$ " wide. Manufactured by R.C.A. Objective, f/0.9 Schmidt system, 2.4" focal length,  $3\frac{1}{4}$ " aperture. Excellent condition, with waterproof carrying case. Price **\$35.00**



**INFRARED FILTERS**

$5\frac{3}{8}$ " diam. x  $\frac{1}{8}$ " thick. Price **\$2.95** each



**BAUSCH & LOMB f/6  
AERO-TESSAR LENS**

Focal length 24". Completely mounted with iris and shutter. Excellent for wide-field telescope. All in excellent condition. Price **\$25.00**



**GOVERNMENT-DESIGNED  
ERECTING EYEPIECE**

The eyepiece system is housed in a precision barrel, with provision for varying the separation of the components and the magnification. The combination consists of a Kellner eyepiece of approximately 25-mm. focal length in tandem with a pair of achromatized doublets. This latter unit has a focal length of about 25 mm. and functions as the erector, or as a symmetrical eyepiece when used alone. The erector system is mounted in a threaded cell which may be turned in or out to vary the spacing and power. The entire assembly has a focal length of about 10 mm., and when used complete or as components yields hard, wire-sharp images, 1-3/16" in O.D. by  $4\frac{3}{4}$ " in length; equipped with a rubber eyeshield. Price **\$5.95**



**8-POWER ELBOW TELESCOPE**

2" objective; Kellner eyepiece; Amici erecting prism; 4 built-in filters; reticle illumination; field is  $6^{\circ}$  (325 ft. at 1,000 yds.). The focusing 28-mm. eyepiece allows focus from 15 ft. to infinity. Weight, 5 lbs. These were used by the government for antiaircraft. Government cost over \$200.00. Price **\$12.50**

**5" SCHMIDT ULTRA-HIGH-  
SPEED OBJECTIVE LENS  
SYSTEM**

Eastman Kodak infrared receiver, formerly known as U. S. Navy Metascope, Type B. 7" long with 5" Schmidt ultra-high-speed objective lens (approx. f/0.5). Elaborate optical system, many coated lenses. Uses two penlight batteries. Cost government approx. \$300.00. Factory-new. Shipping weight 9 lbs. Price **\$19.95**



Waterproof carrying case extra. Shipping weight 3 lbs. Price **\$3.00**


**f/8 40" AERIAL CAMERA LENS**

Mfgd. by Bausch & Lomb. Completely mounted with iris. Original negative size 9" x 9". In excellent condition. Price **\$30.00**



**BINOCULAR EYEPIECE**

Completely assembled with coated lenses. Focal length 22.5 mm.; used on 6 x 30 binoculars. Price **\$3.95**




**16-mm. AERIAL GUN CAMERA,  
MODEL AN-N6**

Takes standard 16-mm. magazine load (50'). Takes 16, 32, or 64 frames per second. Wollensak Type V lens, 35-mm. f/3.5. Excellent condition. Uses 24-volt d.c. Price **\$22.50**



**8-DAY AIRCRAFT CLOCK**

24-hr. dial and civil date indicator. Center sweep second hand. Luminous figures and hands on black face. Case is made of black plastic,  $3\frac{1}{8}$ " mounting. Mfgd. by Waltham Watch Co. Price **\$20.95** postpaid



**SEXTANT #AN5851-1**

Bubble type (with alternate averaging device). Mfgd. by Bendix. Unit contains a horizontal indicator, a precision averager (for two-minute readings), an astigmatizer lens, bubble-level prism, spring-wound motor, etc., with carrying case. Shipping weight 20 lbs. Price **\$12.50**

Send full amount with order. All prices, except as noted, net f.o.b. Pasadena, Calif. No C.O.D.'s, please.

# C & H SALES CO.

2176 EAST COLORADO STREET  
PASADENA 8, CALIFORNIA  
Murray 1-7393

## CHECKING CONCENTRICITY OF MAKUTOV CORRECTING LENSES

**N**UMEROUS jigs have been constructed by amateurs to test for wedge in a Maksutov telescope's correcting lens. However, Warren I. Fillmore, in *Construction of a Maksutov Telescope*, page 9, mentions that only a micrometer and a good eye are needed. I have checked the accuracy of this method and recommend it to the amateur.


Molded blanks must be surfaced before this test can be used. After rough grinding of both sides, a flat lip is put on the front surface, reducing the effective size of the lens. Since the diameter of the blank is usually 6.4" to 6.8", the aperture should be brought down to the design figure of 6". The lip gives a firm mounting surface and avoids having a sharp edge that is liable to chip or flake at the slightest impact. Figuring increases the aperture slightly, but after polishing the lip can be reground to the correct size, which will also remove any turned-down edge.

For accuracy in measuring the wedge, the lip must be of the same width all around the blank. An excellent gauge for this is a vernier caliper reading to 0.001", used with a high-power magnifier.

The wedge-checking procedure, though simple, is necessary even for professionally prepared correctors generated on a diamond-wheel surfacing machine. The

spindle of a 1" micrometer is brought just tangent to the circle formed by the concave surface and the flat lip. To avoid damaging the lens be sure to open the micrometer sufficiently to permit it to be placed over the edge. Turn the spindle gently until it makes contact with the front concave surface while the anvil of the micrometer touches the back curve. Contrary to Mr. Fillmore's experience,

This photograph by Robert E. Cox shows how to place a one-inch micrometer to check the correcting lens of an f/15 Maksutov telescope for wedge effect. At this stage, 0.021 inch had yet to be removed from the thickness to complete the grinding of the lens. Placing the spindle at the very edge of the curve gives a consistent reference position.



I found that setting the micrometer tangent to the edge of the glass is very important when readings are taken with the vernier to the nearest 0.0001".

To be sure that the spindle makes full contact all around its edge with the glass surface, rock the anvil against the convex side slowly and gently toward and away from the center of the lens and in a plane at right angles to this line.

During the checking procedure, I made each micrometer reading first by estimat-

## The Very Popular AMATEUR Observa- Dome



### MODEL NO. 9.167-AG • 9 FT. NOMINAL DIAMETER

Our Amateur dome has recently been redesigned to incorporate double transverse shutters. It is entirely self-supporting and requires no interior ribs, following very closely the same general design as our standard aluminum Observa-Dome. Construction is of 24-gauge galvanized steel and is finished with a protective primer coat.

Our Amateur Observa-Dome is priced at only \$622.00, completely assembled and crated, f.o.b. our plant. Shipping weight, 400 pounds.

One of these latest Amateur Observa-Domes is being shipped to Mr. Octavio Fernandez Reynoso, Paris 15, Mexico City 4, D. F.

P. O. BOX 885  
143-47 WESLEY AVE.  
JACKSON, MISSISSIPPI



ing tenths of a regular division on the spindle, and it was possible to hold the accuracy within 0.0002" without using the vernier. This means that a careful amateur can test with an ordinary 0.001" micrometer and still hold the wedge well under the 0.001" given in the design figures as the maximum permitted deviation from uniformity.

However, makers of Maksutov telescopes have found, in general, that if possible the error should be reduced to 0.0001" or less for surfaces over 4" in diameter. Checks of the blank during surfacing the concave side of the corrector shown in the picture indicated a wedge under 0.002" at all times, so final correction was left for the convex side. After being worked until no wedge could be detected with the vernier micrometer (reading to 0.0001"), the lens was checked on a professional jig, using a 0.0001" dial indicator. No error greater than 0.00013" could be found, showing that a micrometer can be used without elaborate jigs, even for the most discriminating amateur optical worker.

In the last stages of fine grinding, precautions should be taken to prevent the end of the spindle or the anvil from marking the lens surfaces. The metal, since it is especially hardened, can easily scratch the glass.

ROBERT E. COX  
327 S. Main St.  
O'Fallon, Mo.

# FIRST STEP TO FINER PERFORMANCE!

## Easy Low-Cost Way To Enjoy Thrills Of Astrophotography

Convert your present 35-mm. Camera with this new Adapter

Developed by Criterion especially for single-lens reflex 35-mm. cameras with interchangeable lenses. Simply remove present lens mount and add this precision-made achromatic lens that adapts camera to eyepiece of any telescope — refractor or reflector. When ordering, be sure to specify make of your camera.

For 35-mm. cameras with SCREW-TYPE lens mount

Cat. #CPA-36 for 1 1/4" eyepiece holders. \$19.00

Cat. #CPA-37 for 4" standard Dynascopes \$17.50

For 35-mm. cameras with BAYONET-TYPE lens mount

Cat. #BPA-36 for 1 1/4" eyepiece holders. \$22.50

Cat. #BPA-37 for 4" standard Dynascopes \$21.00

New!



**Dyn-o-Astro 35-mm. Camera**

at less than regular price of camera alone!

No need to be an expert photographer to take astrophotos with this single-lens reflex-type, precision-made 35-mm. camera. No complicated settings. No plates or filmholders to load and unload. No worries about missing target. Large focusing screen shows you exactly the viewing field up to moment of shooting.

— No accidental double exposures. Winding knob automatically advances film, positions mirror, winds shutter, counts exposures.

— Dual-speed setting knob controls both fast and slow shutter speeds. Color coding makes mistakes impossible.

— Integrated magnifier for critical focusing even on dim objects.


Takes time exposures and also has speeds up to 1/500 second. Guaranteed for 2 years. Complete, ready for use.

Model CP-35 fits 1 1/4" eyepiece holder \$89.00 postpaid

Model CP-36 fits all standard 4" Dynascopes \$85.00 postpaid

### Dyn-o-Swing Camera Support

fits any 4" or 6" telescope without drilling!



Takes guesswork out of astrophotography by providing precise centering of camera over eyepiece. Smooth rack-and-pinion adjustment. Easily attached or detached from tube, with rigid clamp for maximum stability. Special arm lets you swing camera away from eyepiece for visual observation, with instant return for photographing. Can be used with almost any camera. Postpaid.

Cat. #CS-4 to fit all 4" Dynascopes \$17.50

Cat. #CS-47 to fit all 4 3/4" O.D. tubes \$18.95

Cat. #CS-67 to fit custom 6" Dynascopes and 7 1/4" O.D. tubes \$19.95

Catalog F, describing other accessories and parts, cheerfully sent on request. Satisfaction guaranteed, or money refunded. All items sent postpaid. We pay all postage costs. No shipping, crating, or packing charges. Send check, cash, or money order for immediate delivery.

### Criterion Achromatic Barlow

This precision Barlow features an achromatic, coated lens, perfectly centered and corrected for finer resolution than you've ever experienced. Intensively tested and proven superior, achromatic lens is mounted in a precision cell. Gives 2.4x magnification. Definition is clear and sharp, hard to the very edge.

Model CP-4 fits standard 1 1/4" eyepieces \$17.50

Model CP-2 fits small 0.946" eyepieces \$15.50

### Paraboloidal Mirrors

Aluminized, zircon-quartz overlaid. Guaranteed unconditionally to perform to the limits of resolution. Hand-corrected to exacting specifications, and guaranteed superior.



4" f/8	\$42.00
4" f/10	\$31.00
6" f/8	\$55.00
6" f/9	\$50.00
8" f/8	\$89.00
10" f/7	\$179.00
12 1/2" f/7	\$299.00

All mirrors made of PYREX-brand glass.

A tolerance of 5% in focal length is customary. A deposit of 1/3 is required with orders for 8" to 12 1/2" mirrors.

### Reflecting Telescope Mirror Mounts

Mounting the mirror to your scope correctly is most important. Criterion mounts are especially well designed, and are made of cast aluminum with brass mounting and adjustment screws. One section fits tube, other section holds mirror. Alignment accomplished by three spring-loaded knurled adjusting nuts. Outer cell designed to fit into or over your tube. Sufficient space left between the two cells. All drilled and tapped. Complete with holding clamps, springs, nuts, etc. Ready for use. Will prevent vibration and hold alignment once set. Will hold mirror without distortion of surface figure.



3"	\$3.00	6"	\$6.50
4"	\$3.50	8"	\$12.50
5"	\$4.00	10"	\$19.50

### Complete Eyepieces




Finest quality. They are precision machined, mounted in standard 1 1/4" outside diameter barrels. Can be taken apart for cleaning. Designed to give sharp flat field clear to edge.

Huygens 18-mm. f.l. (1/3")	\$7.50
Kellner 9-mm. f.l. (1/3")	7.90
Kellner 7-mm. f.l. (9/32")	8.50
Kellner 12.7-mm. f.l. (1/2")	9.50
Kellner 18-mm. f.l. (1/2")	9.50
Kellner 30-mm. f.l. (1-3/16")	12.50
Orthoscopic 6-mm. f.l. (1/4")	12.50
Orthoscopic 4-mm. f.l. (5/32")	14.50

### Revolving Turret

The Criterion Revolving Turret holds three eyepieces so that, as desired, the power of the telescope can be changed by merely turning the turret to a different ocular. Click stop insures positive and accurate positioning of each eyepiece. Turret holds eyepieces of standard 1 1/4" outside diameter. Fits into the holder of any refractor or reflector telescope that uses 1 1/4" eyepieces. Requires no alteration or adjustment and can be attached as easily as putting eyepiece into scope. Made of brass and aluminum with polished chrome-plated barrels.

Cat. #SRT-350 \$14.50



### Rack-and-Pinion Eyepiece Mount


The most mechanically perfect focusing is by rack and pinion. This mount takes standard 1 1/4" eyepieces. Full 3 1/2" of travel — more than ever before. Accommodates almost any type of eyepiece — positive and negative. Two knurled focusing knobs, variably tensioned and positioned. Solid cast-metal sliding brass tube — close tolerance prevents looseness. Mount aligns itself to any type tube. Four mounting holes, nuts and bolts included. Eye mount has square-rod-type diagonal holder which prevents loose alignment and vibration. Rod tempered to minimize temperature changes. Adjustable for 3" to 8" scopes, also 12" scopes if so specified at no extra cost. Order one or more of the complete eyepieces described below at the same time you send for this rack-and-pinion device, which accommodates any of our eyepieces perfectly.

Cat. #SU-38 \$7.95 postpaid

### New Model Eyepiece Mount

Same features as above but has wider base that is contoured to match the curve of a 7" to 8" diameter tube. Makes professional appearance. Furnished without Diagonal Rod \$9.95

Diagonal Rod — Cat. #SU-9R \$1.00



### Achromatic Finder Scopes

Two models: 6x, 30-mm., and 10x, 42-mm. Coated achromatic air-spaced objective, cross-hairs, built-in duralumin tube finished in white enamel, dewcap. Sliding focus adjustment. Can also be used as excellent hand telescopes for wide-field views of the sky. Fit Mount Bracket #SF-610.

6 x 30 Achromatic Finder	\$12.50
10 x 42 Achromatic Finder	\$18.00
Mount Bracket #SF-610	\$9.95

### Wide-Angle Erfle Eyepiece

Our 16.3-mm. Erfle wide-angle eyepiece has a 75° field. Astonishing wide-angle views. Coated. Highest precision and specifically designed for telescopic use. Chrome barrel. Guaranteed to be superior in every respect.

Cat. #SE-63 — 1 1/4" O.D. \$18.50

Cat. #SE-62 — 0.946" O.D. \$16.50

### Four-Vane Diagonal Holders

Criterion 4-vane diagonal mountings are fully adjustable, will hold elliptical diagonals in perfect alignment. Made of brass, chemically blackened. Precision adjusting screws center flat and vary its angle so that primary and secondary mirrors can be set in perfect alignment. Thin vanes with special adjustable studs.



Cat. #	Minor-Axis Size	For Tubes	Price
S-51	1.25"	6 1/4" to 7 1/2"	\$10.00
S-52	1.30"	6 1/2" to 7 1/2"	10.00
S-53	1.50"	8 1/2" to 9 1/2"	10.00
S-54	1.75"	9 1/2" to 10 1/4"	12.50
S-55	2.00"	11" to 11 1/2"	14.95
S-56	2.50"	Specify tube I.D.	19.95

## CRITERION MANUFACTURING CO.

Manufacturers of Quality Optical Instruments

Dept. STP-32, 331 Church St., Hartford 1, Conn.



# CELESTIAL CALENDAR

Universal time (UT) is used unless otherwise noted.

## AN EARLY-MORNING OCCULTATION OF ALDEBARAN

**M**ORNING TWILIGHT will be brightening the sky on July 9th as the waning crescent moon occults Alpha Tauri for watchers in the southeastern and south-central United States. The 1st-magnitude star will disappear at the bright edge of the 26-day-old moon, an event that occurs after sunrise in New England and before moonrise in Texas, where predictions are given only for emersion from the moon's dark limb.

At Washington, D. C. (Station C), conditions will permit watching the occultation's beginning. Disappearance there is predicted for 5:33.1 Eastern daylight time, at position angle 27° (measured eastward from the moon's north point). Reappearance at 6:18.1, 299°, will be a difficult telescopic observation requiring excellent sky conditions, as the sun will have risen

half an hour before this, at 5:49 EDT.

But with sunup generally later at places west and south of Washington, the occultation circumstances become more favorable. The moon should be watched as soon as it rises that morning. The gap between it and the star will continually close, and Aldebaran should be visible with telescopic aid right up to the moment of contact with the moon's limb. Reappearance at Station G (Texas) is scheduled for 5:00 Central daylight time, during morning twilight, at position angle 302°.

A grazing occultation is predicted for more northerly observers, for example in Indiana and Illinois. Additional prediction data may be found in the SKY AND TELESCOPE Occultation Supplement (December, 1960, page 349).

## MINOR PLANET PREDICTIONS

Dembowska, 349, 10.1. July 30, 22:12.9 —24:12. August 9, 22:05.3 —24:57; 19, 21:56.5 —25:34; 29, 21:47.7 —25:58. September 8, 21:39.7 —26:05; 18, 21:33.6 —25:54. Opposition on August 20th.

After the asteroid's name are its number and the approximate visual magnitude expected at opposition. At 10-day intervals are given its right ascension and declination (1950.0) for 0<sup>h</sup> Universal time. In each case the motion of the asteroid is retrograde. Data are supplied by the IAU Minor Planet Center at the University of Cincinnati Observatory.

## MOON PHASES AND DISTANCES

Last quarter	July 5, 3:33
New moon	July 12, 19:12
First quarter	July 20, 23:14
Full moon	July 27, 19:51
Last quarter	August 3, 11:48
<b>July</b>	
Apogee 15, 11 <sup>h</sup>	252,400 mi. 29° 25"
Perigee 28, 9 <sup>h</sup>	222,200 mi. 33° 25"
<b>August</b>	
Apogee 11, 17 <sup>h</sup>	252,600 mi. 29° 24"

## MINIMA OF ALGOL

July 3, 11:55; 6, 8:44; 9, 5:32; 12, 2:21; 14, 23:10; 17, 19:59; 20, 16:48; 23, 13:36; 26, 10:25; 29, 7:14.

August 1, 4:02; 4, 0:51; 6, 21:40.

These minima predictions for Algol are based on a recent timing by D. Engelkeir, and an assumed period of 2.8674 days. The times given are geocentric; they can be compared directly with observed times of least brightness.


## Sky and Telescope Binders

Dark blue fabrikoid binders priced at \$3.50 each, postpaid in the United States; \$4.00 in Canada. Two sizes: Binder C is for volumes up to XVIII; Binder D is for volume XIX and after. If it is desired to put two of the current volumes (a total of 12 issues) into a single binder, order Binder C instead of D. Your name can be gold-stamped for 75¢ extra, each volume number for 50¢; name and volume number, \$1.20; name and two volume numbers, \$1.70. Print all desired lettering clearly.

(Sorry, no orders for binders can be accepted from other countries.)

Sky Publishing Corporation

Harvard Observatory, Cambridge 38, Mass.



This mean light curve for X Ophiuchi was derived from observations made by members of the American Association of Variable Star Observers.

## X OPHIUCHI

**N**EAR the point where the summer constellations Ophiuchus, Aquila, and Hercules meet is the Mira-type variable star X Ophiuchi, at right ascension 18<sup>h</sup> 36<sup>m</sup>.0, declination +8° 42' (1950 coordinates). Maximum brightness of about magnitude 6.8 is forecast for around July 9th this year. At no stage in its 334-day cycle does this object fade out of reach of a 3-inch telescope, minimum light being about 8.6. Ever since its variability was detected by the English amateur T. E. Espin in 1886, X Ophiuchi has been a favorite with visual observers.

The light changes have certain marked peculiarities. Minima are comparatively shallow and flat, and differ very little from cycle to cycle, although, like other Mira stars, the maxima are quite varied in height. These facts suggest that X Ophiuchi is a close double star, whose variable component is sometimes much brighter, at other times fainter than its constant companion. Thus at minimum the latter star would be taken for the variable.

This is actually the case, for W. J. Hussey with the Lick Observatory 36-inch refractor found a 9th-magnitude companion only 0.3 second south of the

variable. Later micrometer measurements have established very slow orbital motion in this visual binary. At Mount Wilson, P. W. Merrill ascertained that the variable is of spectral class M6e, the constant companion of class K0. Hence, X Ophiuchi, as seen unresolved in amateur telescopes, should appear red at maximum and yellow at minimum.

## JUPITER WITHOUT SATELLITES

An unusual event takes place on September 27th, when Jupiter will appear without its bright satellites — all four being either in occultation, eclipse, or transit. This will occur when Jupiter is below the horizon for American observers, but the sequence of events will be visible from Japan and Australia. The following timetable is given by the *American Ephemeris*.

At 13:30 Universal time all four moons will be visible, I to the west of Jupiter, II, III, and IV to the east. At 13:56, III begins a transit across the face of the planet, and II transits at 15:16. The two remaining satellites will vanish nearly simultaneously, I going behind the west limb of the planet at 15:59, IV entering eclipse in Jupiter's shadow at 16:00.

Jupiter will seem to have no moons until 17:30, when III passes off the disk, ending its transit, II doing the same at 18:05. Next, I emerges from eclipse at 19:29, and IV similarly reappears at 20:40.

Jupiter last appeared moonless on September 20, 1949, and will do so again on June 27, 1966. Remarkably, on April 9, 1980, the phenomenon will occur twice in one night, satellite III emerging from occultation and promptly vanishing again into eclipse while the others remain invisible. For a list of occasions in the 19th and 20th centuries when Jupiter is without visible satellites, see the *Journal of the British Astronomical Association* for November, 1931, page 12.


## JULY METEORS

The moon, which is full on July 27th, will interfere with observations of the Delta Aquarid meteors, at maximum two days later. Under more favorable conditions, a single observer might see about 20 meteors per hour at the peak of the 20-day shower. On the 29th the radiant will be just west of Delta Aquarii, at right ascension 22<sup>h</sup> 36<sup>m</sup>, declination -17°; this point will move east and slightly north almost a degree per day.

WILLIAM H. GLENN

## UNIVERSAL TIME (UT)

TIMES given in Celestial Calendar are Universal time (Greenwich civil time) unless otherwise noted. This is 24-hour time, from midnight to midnight; times greater than 12:00 are p.m. Subtract the following hours to convert to standard times in the United States: EST, 5; CST, 6; MST, 7; PST, 8. To obtain daylight saving time subtract 4, 5, 6, or 7 hours respectively. If necessary, add 24 hours to the UT before subtracting, in which case the result is your standard time on the day preceding the Greenwich date shown. For example, 6:15 UT on the 15th of the month corresponds to 1:15 a.m. EST on the 15th and to 10:15 p.m. PST on the 14th.



### THE SUN, MOON, AND PLANETS THIS MONTH

The sun, on the ecliptic, is shown for the beginning and end of the month. The moon's symbols give its phase roughly, with the date marked alongside. Each planet is located for the middle of the month or for other dates shown. All positions are for 0<sup>h</sup> Universal time on the respective dates.

Mercury is stationary in right ascension at 19<sup>h</sup> Universal time on July 8th, before beginning eastward (direct) motion among the stars. On the 19th it reaches greatest western elongation, 20° from the sun, rising about 1½ hours before it. Mercury will be about as bright as Capella, then some 30° northwest of the planet. For a week before and after elongation, Mercury will be far enough away from the sun to be seen. Its disk will be near quarter phase, which will be reached on the 23rd.

Venus is the most brilliant object in the morning sky, of magnitude -3.7 on the 15th, about 100 times as bright as 1st-magnitude Aldebaran, which will be 3° south of the planet as the latter passes through the Hyades during the middle of July. Venus' disk appears in a telescope slightly gibbous, at midmonth 62 per cent illuminated and 18".6 in diameter.

Mars is an inconspicuous object in Leo this month, of magnitude +1.8 and with a disk only 4".4 in diameter on the 15th. It may be seen very low in the west as evening falls.

Jupiter reaches opposition to the sun on the 25th, 381 million miles from the earth. It rises in Capricornus just before sunset, and is visible the whole night as

### VARIABLE STAR MAXIMA

July 4, RU Cygni, 213753, 8.0; 9, T Hydrae, 085008, 7.8; 9, X Ophiuchi, 183-308, 6.8; 12, V Cancri, 081617, 7.9; 24, T Aquarii, 204405, 7.7; 29, R Octantis, 055686, 7.9; 29, R Pegasi, 230110, 7.8.

August 6, T Eridani, 035124, 8.0; 9, W Ceti, 235715, 7.6.

These predictions of variable star maxima are by the AAVSO. Stars are listed only if brighter than magnitude 8.0 at an average maximum. Some, but not all of them, are nearly as bright as maximum two or three weeks before and after the dates for their maxima. The data given include, in order, the day of the month near which the maximum should occur, the star name, the star designation number, which gives the rough right ascension (first four figures) and declination (bold face if southern), and the predicted visual magnitude.

an object of magnitude -2.3 in the southern sky. Telescopically, its disk is 48".1 in equatorial diameter and 44".9 in polar.

Saturn, in Sagittarius about 6° west of Jupiter, reaches opposition on July 19th, 837 million miles from Earth, when its magnitude will be +0.3. The planet's slightly flattened disk has an equatorial diameter of 18".5 and a polar one of

### JUPITER'S SATELLITES

The four curving lines represent Jupiter's four bright (Galilean) satellites: I, Io; II, Europa; III, Ganymede; IV, Callisto. The location of the planet's disk is indicated by the pairs of vertical lines. When a satellite passes in front of Jupiter, its curve crosses the lines. If a moon is invisible because it is occulted by Jupiter or is in its shadow, the curve is broken.


For successive dates, the horizontal lines mark 0<sup>h</sup> Universal time, or 7 p.m. Eastern standard time (4 p.m. Pacific standard time) on the preceding day. Along the vertical scale, 1/16 inch is about seven hours. In this chart, west is to the left, as in an inverting telescope for a Northern Hemisphere observer. At the bottom, "d" is the point of disappearance of a satellite in Jupiter's shadow; "r" is the point of reappearance. From the "American Ephemeris and Nautical Almanac."


16".6. The moon will pass 3° north of Saturn at 7<sup>h</sup> UT on July 27th.

Uranus is in Leo, poorly placed for observation this month, for it sets at the end of evening twilight.

Neptune is an 8th-magnitude object in westernmost Libra, with 1950 co-ordinates 14<sup>h</sup> 26<sup>m</sup>.4, -12° 37' on the 15th. It crosses the meridian before sundown and sets in the southwest before local midnight. The planet's tiny greenish disk is 2".4 in diameter. On the 21st, Neptune becomes stationary in right ascension, as it resumes eastward motion in the heavens.

WILLIAM H. GLENN





## *The race to space may be won in your own school with a Tinsley Telescope*

There are uncountable universes waiting to know the mastery of the Earthborn scientist. Perhaps he will be the dedicated scholar in the first row of your class, or the boy just behind him to the right. The Tinsley 12-inch Telescope will open the secrets of galaxies upon galaxies to them, and to all the young minds who must absorb — and learn to live with — concepts unimaginable to their grandparents.

**THE TINSLEY 12-INCH TELESCOPE** is the finest manufactured in its size range. It is produced with the same painstaking care and engineering Tinsley brings to custom telescopes for universities and for private and government research institutions.

**PRECISE CONTROLS FOR CLASSROOM USE:** The Tinsley 12-inch responds smoothly, as quickly or slowly as you want it to, with perfect freedom and balance. It moves on big angular contact ball bearings of a kind not found in lesser telescopes. Once set, the Tinsley 12-inch tracks accurately without readjustment. Its positive-action clamp mechanism assures this. Motor driven in right ascension to a precise sidereal rate. Additional fast drive provided. Manual declination with antibacklash drive.

**OPTICAL SYSTEM:** Cassegrain, with primary mirror focal length 48"; secondary amplification 4; total effective focal length 192". Four standard eyepieces give a range of magnification from 48 to 384 diameters. Rugged rack-and-pinion focusing assembly has positive locking clamp, and may be removed and replaced with a camera or spectrograph. Optical surfaces will satisfy the Dawes resolution limit. Equipped with a finder of 10 power. Photographic attachments available.


All parts of the Tinsley Telescope have been treated with or made from corrosion-resistant materials • Mounted on 60-inch-high heavy steel pedestal • Accessory compartment • Dustproof lens cover • Can be roof-installed • Plans for a simple, weather-and-vandal-proof shelter available.

☐ \$4,850.00, F.O.B. BERKELEY ☐

For a complete specification folder, write to Department M, Tinsley Laboratories, Inc., 2448 Sixth Street, Berkeley 10, Calif.







## STARS FOR JULY

The sky as seen from latitudes 30° to 50° north, at 9 p.m. and 8 p.m., local time, on the 7th and 23rd of July, re-

spectively; also, at 7 p.m. on August 7th. For other dates, add or subtract ½ hour per week.

The forepart of Serpens, the great snake grasped by Ophiuchus, is just past the

meridian at chart time. Its head is marked by an equiarmed "X" of five stars; from there the constellation may be traced through Alpha, a yellow 3rd-magnitude star in the snake's neck.

# OPTICS

## Special Coated Objectives

BIG 2 1/4" DIAM. — 40" F.L. — \$6.00

These tested achromatic lenses are the same high quality as our "Big Lenses," except for being seconds because of slight edge chips or small scratches.

Cat. No. S1804 ..... Only \$6.00 ppd.

## Mounted Eyepieces

The buying of a lifetime at a great saving. First choice of the amateur and professional astronomer. Perfect lenses set in black anodized standard 1 1/4" O.D. aluminum mounts.

Cat. No.	F.L.	Type	ppd.
S2192	6 mm. (1/4")	Orthoscopic	\$13.25
S1851	6 mm. (1/4")	Ramsden	4.75
S1853	12.5 mm. (1/2")	Ramsden	4.50
S1207	12.5 mm. (1/2")	Symmetrical	6.00
S1251	16 mm. (5/8")	Erfle (wide-angle)	12.50
S1257	16 mm. (5/8")	Orthoscopic	12.50
S1209	18 mm. (3/4")	Symmetrical	6.00
S1211	22 mm. (7/8")	Kellner	6.00
S1835	27 mm. (1 1/16")	Kellner	4.50
S1355	32 mm. (1 1/4")	Plossl	12.50
S1253	35 mm. (1 3/16")	Symmetrical	8.00
S1255	55 mm. (2 1/16")	Kellner	6.00
S1485	56 mm. (2 1/4")	Symmetrical	6.00

COATED LENSES 75 cents extra.

## "Giant" Wide-Angle Eyepiece

COATED LENSES 75 cents extra.

Erfle eyepiece best for low-power performance. Superior six-element design. 65° field, 1 1/2" E.F.L., clear aperture 2 1/8", focusing mount and diopter scale. Combine this eyepiece with our 5" objective No. S1475 for a tremendous light-gathering Rich-Field Telescope.

Cat. No. S1405 (illustrated) ..... \$12.50 ppd.  
Cat. No. S1858 Same as above without diopter scale ..... 9.95 ppd.

● FREE RUBBER EYESHIELD with S1405

Cat. No. S1594 1 1/4"-diam. ADAPTER for eyepiece above ..... \$ 3.95 ppd.

**ADAPTER PLATE AND TUBE.** Aluminum tapered tube and adapter plate, black anodized and finished in black crackle. Slides into our 5" O.D. Aluminum Tubing No. S1311. These parts are used with the eyepieces above to make a Rich-Field Telescope. Details in our catalogue.

Cat. No. S2300 ..... \$18.00



## Wide-Angle Erfle

Brand-new eyepiece with 68° field; coated. E.F.L. 1 1/4". Focusing mount, 3 perfect achromats, 1-13/16" aperture.

Cat. No. S1020 ..... \$13.50 ppd.  
Cat. No. S1593 1 1/4"-diam. ADAPTER for eyepiece above ..... 3.95 ppd.

## An Economical Eyepiece

This mounted eyepiece has two magnesium-fluoride-coated achromatic lenses 29 mm. in diameter. Excellent definition. E.F.L. 1 1/4". Cell fits 1 1/4" tubing.

Cat. No. S1911 Coated ..... \$5.90 ppd.  
Cat. No. S1991 Not Coated ..... 5.25 ppd.

## Rack-&-Pinion Eyepiece Mounts



Here is a wonderful opportunity for you to own a most mechanically perfect Rack-&-Pinion Focusing Eyepiece Mount with variable tension and adjustment. Will accommodate a standard 1 1/4" eyepiece, positive or negative. The body casting is made of lightweight aluminum with black-crackle paint finish. Focusing tube of chrome-plated brass. Focusing tube for reflectors has a travel of 4", for reflectors 2". Mount will fit all size tubing.

### REFRACTOR TYPE


Cat. No. S1494 for 2 1/8" I.D. Tubing ..... \$12.95 ppd.  
Cat. No. S1495 for 3 1/4" I.D. Tubing ..... 12.95 ppd.  
Cat. No. S1496 for 4 3/8" I.D. Tubing ..... 12.95 ppd.

### REFLECTOR TYPE

Cat. No. S1976 (less diagonal holder) ..... \$8.50 ppd.  
Cat. No. S1982 DIAGONAL HOLDER ..... 1.00 ppd.

## 8-Power Elbow Telescope

This telescope has a brilliant image 48° apparent field; actually 325 feet at 1,000 yards. Can be adjusted for focusing 15 feet to infinity. 2" achromatic objective, eyepiece 28-mm. f.l., Amici erecting system. Turret-mounted filters: clear, red, amber, and neutral. Lamp housing to illuminate reticle for night use. Truly the biggest bargain you were ever offered. Original Gov't. cost \$200.




8 x 50

Cat. No. S1686 Not Coated ..... \$13.50 ppd.  
Cat. No. S1975 Coated ..... 17.50 ppd.

## Star Diagonals

Guaranteed to make viewing the heavens with a refractor telescope 100% easier for the observer. Stop needless crouching forever. Choose the STAR DIAGONAL that best suits your requirements. Slides into standard 1 1/4" eyepiece mounts, and aluminum housing, attractive black-crackle finish.



Chromium tubes and

## AMICI STAR DIAGONAL WITH ERECT IMAGE

For comfortable viewing of terrestrial or astronomical objects. The Amici prism gives an erect image.

Cat. No. S2095 ..... \$12.00 ppd.

## RIGHT-ANGLE PRISM STAR DIAGONAL

Contains a high-quality aluminized right-angle prism, wide field. Practical for astronomical use.

Cat. No. S2096 ..... \$12.00 ppd.

## MIRROR STAR DIAGONAL

Diagonal houses an excellent first-surface mirror, flat to within 1/4 wave. Important accessory for refractor telescopes.

Cat. No. S2097 ..... \$12.00 ppd.

## 7 x 30 Wide-Field Finderscope



Newly designed wide-field finderscope for reflector or refractor. Featured in this handsomely conceived black anodized aluminum tube are a 30-mm. coated achromatic objective, a wide-field Kellner eyepiece, and the all-important true crossline reticle for fast, exact locating. Compact and lightweight, 9" long, 1 1/2" diam., 9 oz. May also be used as a hand-held telescope, offering excellent extra-wide-field views of the sky.

Cat. No. S2193 ..... \$9.75 ppd.

## Finder Ring Mounts

Made of cast aluminum with black crackle finish. Each ring has three locking wing screws for adjusting. Base has two holes for mounting screws, and fits any diameter tubing. Easy to attach. Ring mount No. S1963 will accommodate finder above.

Cat. No. S1963 .. 1-15/16" I.D. .. Per Pair, \$3.95 ppd.  
Cat. No. S1964 .. 2-11/16" I.D. .. Per Pair, 3.95 ppd.

## New Prismatic Telescope

Coated



80-mm. Objective

Big 80-mm.-diam. (3 1/4") objective, for spotting or astronomical use. Will show stars of the 11th magnitude — 100 times fainter than the faintest visible to the naked eye. Table-top tripod included, extended 18", closed 13 1/4". Length of scope 16 1/4". Combined weight of scope and tripod is 5 1/4 lbs. All coated optics.

### FIVE EYEPIECES

Power	Field at 1,000 yards	Exit pupil diam.	Relative Brightness
15x	122 ft.	5.4 mm.	20
20	81	4.0	16
30	61	2.7	7
40	49	2.0	4
60	32	1.3	1

Cat. No. S2052 ..... \$59.50 ppd.


60-mm.-diam. Scope. Same as above but with smaller objective. Equipped with same five eyepieces — 15x, 20x, 30x, 40x, 60x. Coated. With tripod.

Cat. No. S1721 ..... \$42.95 ppd.

## 4" Reflector Telescope

45x-225x

4x Finder — Wood Tripod



Study the sun and its spots. Observe galaxies, clusters, nebulae, and the Milky Way. Enjoy the thrill of being able to pick out small craters on the moon and see detail in the larger craters. A beautiful lunar landscape is brought to your doorstep. Bring distant events close to you; witness nature in action — eclipses, heavenly objects, and orbiting satellites. A high-quality telescope with an f/9 mirror. It has three eyepieces, 4-mm. (225x), 6-mm. (150x), 20-mm. (45x), and a free Barlow lens which will double the power. This will give 90x with the 20-mm. eyepiece, and 300x with the 6-mm. if observing conditions permit. Equipment tray keeps eyepieces clean and at hand. Sun and moon filters are included. Collapsible wood tripod makes scope portable, gives it stability and compactness. Finished in gray enamel with black trim. Imported. Shipping weight about 20 lbs.

FREE BARLOW LENS PLUS FREE BOOK "DISCOVER THE STARS" will come with scope.

Cat. No. S2273 ..... f.o.b. Lynbrook, N. Y., \$59.50

## COATED BINOCULARS

Beautifully styled imported binoculars. Precision made. LOW, LOW PRICES!



American Type



"Zeiss" Type

Complete with carrying case and straps. American type offers a superior one-piece frame and clean design. Individual Focus (IF) or Center Focus (CF).

Cat. No.	Size	Field at 1,000 yards	Type	Price, ppd.
S1149	6 x 15 IF	360 ft.	Opera	\$12.75
S1436	6 x 30 CF	395	"Zeiss"	18.75
S1435	6 x 30 IF	395	"Zeiss"	16.75
S1438	7 x 35 CF	341	"Zeiss"	20.75
S1437	7 x 35 IF	341	"Zeiss"	17.95
S1771	7 x 35 CF	341	American	23.50
S1439	7 x 35 CF	578	American*	35.00
S2191	7 x 50 CF	530	American*	42.50
S1106	7 x 50 CF	372	"Zeiss"	24.95
S-961	7 x 50 IF	372	"Zeiss"	22.50
S1503	7 x 50 CF	372	American	32.50
S1443	8 x 30 CF	393	"Zeiss"	21.00
S1229	8 x 30 IF	393	"Zeiss"	18.25
S1108	10 x 50 IF	275	"Zeiss"	26.75
S1442	20 x 50 CF	183	"Zeiss"	33.75

\*Wide-angle 11°

\*\*Wide-angle

Add 10% Federal tax to all prices above.



## Monoculars

Brand new, coated optics, complete with pigskin case and neck straps.

Cat. No. S1576	6 x 30	\$10.00 ppd.
Cat. No. S1577	8 x 30	11.25 ppd.
Cat. No. S1578	7 x 35	12.50 ppd.
Cat. No. S1303	7 x 50	14.75 ppd.
Cat. No. S1579	16 x 50	17.50 ppd.
Cat. No. S1580	20 x 50	20.00 ppd.

## New! Zoom Binoculars

Six different powers at your fingertips in the same binocular: 7x, 8x, 9x, 10x, 11x, and 12x. Positive fingertip control accomplished by rotation of eyecups. Popular Center Focus, American type, 40-mm. objectives, all coated optics. Complete with carrying case and straps. Add 10% Federal tax.

Cat. No. S2303 ..... \$39.75 ppd.

## FREE CATALOG

Millions of lenses, prisms, eyepieces, rack-and-pinion eyepiece mounts, cells, aluminum tubing, mirrors, binoculars, telescopes, parts, accessories, and so forth. Write today!

We pay POSTAGE IN U. S. — C.O.D.'s you pay postage. Satisfaction guaranteed or money refunded in 30 days.

# A. JAEGER'S

6915 MERRICK RD., LYNBROOK, N. Y.

## You'll find these fully professional features in every Custom DYNASCOPE®!

Exquisite optical systems, fully achromatic and corrected to meet the most critical research standards • Massive equatorial mounts that provide exceptional stability and embody the latest mechanical refinements for ease of operation. • Full sets of matched orthoscopic and achromatic eyepieces with wide flat fields sharp to extreme edges. • Needle bearings carefully fitted on substantial shafts to assure movements so fine that they equal the ease and precision of a fine watch. • Lifetime materials throughout, for atmospheric resistance and dimensional stability. • Fully rotating tubes, with counterpoise to provide perfect balance during rota-

tion. • Rotary secondary supports to minimize diffraction. • Exclusive Dyna-mesh rack and pinion focus for smoother, more accurate focusing action. • Two-way pier tripods (Pat. Pending) in 6-inch and 8-inch models. 45-pound pier can be mounted permanently. Portable tripod is instantly removable for field trips. • Highly accurate electric drives, free from backlash, with fully compensating clutch. • 8 x 50 coated achromatic finderscopes with crosshairs and extra-large 6° fields. • 7"-diameter setting circles spoked for easy setting. Graduated with verniers. • Super sensitive slow motion flex line control • Adjustable to all latitudes.

## MASTERPIECES *of precision performance!*



*All sizes of these superb instruments are available for unusually prompt delivery . . . including this Deluxe 10-inch model*

## CUSTOM DYNASCOPES®

in your  
choice of  
6"-8"-10"  
12.5"-16"  
models

Every Custom DYNASCOPE is designed and built to meet such high standards and exacting specifications that it is the overwhelming choice of schools, colleges, universities, and many well-known professional astronomers. Each of these magnificent instruments incorporates features that are years ahead in optical engineering, yet based on time-tested principles. Each is painstakingly produced and assembled by craftsmen who put careful workmanship before

speed. Each is required to pass, flawlessly, a long series of rigid performance and operational tests. Nothing is overlooked to make sure that these superb telescopes are the finest in their class, regardless of price. Whichever size you select, we guarantee that it *must* measure up to this standard or your purchase price will be refunded in full. You can order with absolute confidence . . . and with the exciting expectation of new thrills and experiences ahead.



**Colorful  
New Brochure  
Yours FREE!**

Send today for this fascinating new booklet that gives you the full specifications on all Custom Dynascopes, plus helpful information to guide you in the selection of your next telescope. A postcard will bring your copy FREE!

### CUSTOM DYNASCOPES — PRICE AND DELIVERY SCHEDULE

CONVENIENT TIME PAYMENT PLANS AVAILABLE ON ALL MODELS

SIX-INCH		EIGHT-INCH		10-INCH	12.5-INCH	16-INCH
STANDARD	DELUXE	STANDARD	DELUXE	DELUXE	DELUXE	OBSERVATORY MODEL
\$265 Complete	\$475 Complete	\$385 Complete	\$595 Complete	\$895 Complete	\$1,265 Complete	\$4,200 up
Shipping Weight 80 lbs.	Shipping Weight 150 lbs.	Shipping Weight 145 lbs.	Shipping Weight 185 lbs.	Shipping Weight 245 lbs.	Shipping Weight 300 lbs.	Equipped To Your Specifications
21 Day Delivery		21 Day Delivery		30 Day Delivery	30 Day Delivery	

Standard Six and Eight Inch Models Do Not Include Electric Clock Drive, Setting Circles, Slow Motion Control Or Pier  
All Telescopes Are Shipped F.O.B. Hartford, Connecticut • No Additional Charge For Packing and Crating

© Copyright 1960 Criterion Mfg. Co.



® TM Registered U.S. Pat. Office

**CRITERION MANUFACTURING CO.** *Manufacturers of Quality Optical Instruments*  
**DEPT. D-51, 331 CHURCH ST., HARTFORD 1, CONN.**

PRINTED BY WELLESLEY PRESS, INC.



Johnny isn't watching the late show tonight...



The shadows that shiver and shake on the TV screen are shivering and shaking in somebody else's living room tonight. Johnny has discovered something new.

He's traded the fleeting, flickering "thrills" of the 24 inch screen for the timeless excitement and majesty of the night sky.

He's traded the nervous rattle of the private eye's gun for a ringside seat at the stupendous nightly fireworks in the heavens.

He has, in short, discovered astronomy.

Nothing better could happen than what happened to Johnny. And it happened simply because someone took the trouble to awaken, nourish and satisfy a lifetime of curiosity in Johnny by making him the gift of a fine telescope.

Someone, not so long ago, gave Johnny a Unitron.

Johnny abandoned his 24" screen for this 2.4" Unitron refractor, complete with its handy, handsome, easily portable carrying case.



This is a close-up of the Unihex Johnny is using. It's 6 eyepieces in one, an exclusive with Unitron. One of a complete line of accessories to multiply your viewing pleasure.

**UNITRON**

INSTRUMENT COMPANY • TELESCOPE SALES DIV. — 66 NEEDHAM ST., NEWTON HIGHLANDS 61, MASS.

See pages 36 and 37.

